



ENHANCED BARRIER PRECAUTIONS (LTCFs)



PRECAUCIONES CON BARRERAS REFORZADAS (CENTROS DE LARGA ESTANCIA)

Family/Visitors should not visit if having signs or symptoms of an infection or a communicable disease. Visitation also based on facility's policy.

Los familiares y visitantes no deben visitar si tienen señales o síntomas de infección o de una enfermedad contagiosa. Las visitas también dependen de la política de la instalación.

Follow instructions below before entering room.

Antes de entrar a la habitación, siga las instrucciones a continuación.



Everyone must:

Todos deben:

Clean hands before entering and after leaving room.

Lavarse las manos antes de entrar y antes de salir de la habitación.

All Healthcare Personnel must:

Todo el personal de atención médica debe:

Wear gloves and gown for the following High-Contact Resident Care Activities:

- Dressing Bathing/Showering
 - Transferring
 - Changing Linens
 - Providing Hygiene
- Changing briefs or assisting with toileting
- Device care or use: central line, urinary catheter, feeding tube, tracheostomy
- Wound Care: any skin opening requiring a dressing

Usar guantes y bata para las siguientes actividades de alto contacto durante el cuidado de residentes:

- Vestir, bañar, duchar, trasladar, cambiar la ropa de cama.
- Proporcionar higiene, cambiar la ropa interior o ayudar a usar el baño.
- Cuidado o uso de dispositivos: vía central, sonda urinaria, sonda de alimentación, traqueostomía.

Additional PPE may be required per Standard Precautions.

Es posible que se exija utilizar equipo de protección personal adicional según las precauciones estándar.

Enhanced Barrier Precautions

Not intended for acute care or long-term acute care (LTACs)

All residents with any of the following:

- Infection or colonization with a MDRO when Contact Precautions do not apply. At a minimum include resistant organisms targeted by CDC (e.g., Pan-resistant organisms, Carbapenemase producing *Enterobacteriales*, Carbapenemase producing *Pseudomonas* spp., Carbapenemase-producing *Acinetobacter baumannii* and *Candida auris*)
- Wounds (skin opening that requires a dressing) and/or indwelling medical devices (e.g., central line, urinary catheter, feeding tube, tracheostomy/ventilator) regardless of MDRO colonization status

Facilities may consider applying Enhanced Barrier Precautions to residents infected or colonized with other epidemiologically-important MDROs based on facility policy.

Residents are not restricted to their rooms or limited from participating in group activities

Perform Hand Hygiene

- Alcohol based handrub or wash with soap and water if visibly soiled before and after leaving the room.

Personal Protective Equipment

Wear gloves and a gown for the following High-Contact Resident Care Activities.

Do not wear the same gown and gloves for the care of more than one resident.

High-Contact Resident Care Activities include:

- Dressing
- Bathing/Showering
- Transferring
- Changing Linens
- Providing Hygiene
- Changing briefs or assisting with toileting
- Device care or use: central line, urinary catheter, feeding tube, tracheostomy
 - Wound Care: any skin opening requiring a dressing

Take off and dispose in this order

- Gloves– Carefully remove to prevent contamination of hands upon removal
- Gown– Carefully remove to prevent contamination on clothing or skin
- Alcohol based handrub or wash with soap and water if visibly soiled

Dishes/Utensils:

No special precautions. Should be managed in accordance with routine procedures.

Appropriate Use, Cleaning, and Disinfection of medical equipment and devices

Follow facility policies and procedures for cleaning and disinfection of medical equipment and devices. All shared equipment must be cleaned between patients.

Linen and Textile Management:

Bag linen and trash in patient/resident room (double bagging of trash or linen is not necessary unless outside of bag visibly contaminated). Handle linen and textiles carefully.

Duration of Precautions

Intended to be in place for the duration of a resident's stay in the facility or until resolution of the wound or discontinuation of the indwelling medical device that placed them at risk