

Isolation Precautions

Attachment 2: Quick Glance for Respiratory Panel Isolation Precautions*

| Respiratory Panel | Precautions | | |
|---|-------------|--|---|
| | Type* | Duration † | Comments |
| Influenza A Influenza B Adults Pediatric and immunocompromised persons of all ages | D | 7 days adults 10 days for pediatric and/or immunocompromised patients | Droplet Precautions. Single patient room when available or cohort; avoid placement with high-risk patients; mask patient when transported out of room. |
| Adenovirus | D, C | DI | Precautions until no longer symptomatic or returned to baseline respiratory status |
| Metapneumovirus | D,C | DI | Precautions until no longer symptomatic or returned to baseline respiratory status |
| Parainfluenza 1 Parainfluenza 2 Parainfluenza 3 Parainfluenza 4 | D, C | DI | Precautions until no longer symptomatic or returned to baseline respiratory status. |
| RSV | D, C | DI | Precautions until no longer symptomatic or returned to baseline respiratory status, except in NCCC. Infants in NCCC will remain on precautions for 1 month after positive test. |
| Rhino/Entero | D, C | DI | Precautions until no longer symptomatic or returned to baseline respiratory status |
| Coronavirus (not SARS) | D, C | DI | Precautions until no longer symptomatic or returned to baseline respiratory status |
| Coronavirus (SARS-CoV2) | SAC | See link | Per Discontinuation of Isolation Criteria for COVID Patients protocol |
| <i>Mycoplasma pneumoniae</i> | D | DI | Precautions until no longer symptomatic or returned to baseline respiratory status |

Isolation Precautions

| | | | |
|---------------------------------|---|-----------------------------------|--|
| <i>Chlamydia pneumoniae</i> | D | DI | Precautions until no longer symptomatic or returned to baseline respiratory status |
| <i>Bordetella pertussis</i> | D | For 5 days of appropriate therapy | Precautions until patient has completed 5 days of appropriate therapy |
| <i>Bordetella parapertussis</i> | D | For 5 days of appropriate therapy | Precautions until patient has completed 5 days of appropriate therapy |

***When a Respiratory Viral/Pathogen Panel is sent the patient should be placed on Droplet plus Contact Precautions until test finalized and then follow precautions as listed above.**

DI = Duration of Illness (until symptoms have resolved)

D = droplet precautions.

C = contact precautions.

S = standard precautions.

SAC = special airborne and contact precautions