



Definitions and Surveillance for Healthcare Associated Infections (HAIs) in Long-term Care

Evelyn Cook, RN, CIC
Associate Director SPICE



1

How confident are you that your facility has a strong infection prevention program that includes all the necessary elements?

- A. Completely confident
- B. Somewhat confident
- C. Not confident
- D. Have NO idea



2

Do you believe you have the skills and the qualifications to oversee the infection prevention program?

- A. Yes
- B. No
- C. No way; No how

SPICE 

3

If you wanted to compare your IP surveillance data to another NH in your community that cared for a similar resident population, how confident are you that events will be tracked the same way?

- A. Very confident
- B. Slightly confident
- C. Not confident at all
- D. Not sure if I can compare my own data from one year to the next

SPICE 

4

What standardized definition does your facility use for surveillance?

- A. National Healthcare Safety Network (NHSN)
- B. Revised McGeer Definitions
- C. Loeb Criteria
- D. When the physician documents an infection
- E. No standardized criteria
- F. A and B



5

DISCUSS**DESCRIBE****DISCUSS**

The
importance
of
surveillance

Standardized
surveillance
definitions

Ways to
implement
and apply
surveillance
definitions



6

- “Surveillance is a comprehensive method of measuring outcomes and related processes of care, analyzing the data, and providing information to members of the healthcare team to assist in improving those outcomes and processes (APIC Text)
- “Surveillance system must include “**routine, ongoing, and systematic collection, analysis, interpretation, and dissemination** of surveillance data to identify infections (i.e., HAI and communicable-acquired), infection risks, communicable disease outbreaks and to maintain or improve resident health status:” (CMS 8/24)

SPICE 

7

Rationale for Conducting Surveillance



- One of the most important aspects of an IP’s responsibilities
- Should cover residents, staff, contractors (in the facility) and visitors
- Include process and outcome measures

*Establish Baseline Data
Reduce Infection Rates
Detection of Outbreaks
Monitor Effectiveness of Interventions
Education of HCP*

Required as a Component of Plan

SPICE 

8

Types of Surveillance

- Total (or Whole) House Surveillance
- Targeted Surveillance
- Combination Surveillance Strategy



SPICE 

9

Total (Whole House)

- Monitor:
 - All infections
 - Entire population
 - All units



| Pros | Cons |
|---------------------------|--|
| Monitor all infections | Overall rate not sensitive or risk-adjusted |
| Include entire population | No trends or comparison |
| | Labor intense and inefficient use of resources |
| | Not based on risk assessment |

SPICE 

10

Priority Directed (Targeted)

- Focus on:
 - Care units
 - Infections related to devices
 - Invasive procedures
 - Significant organisms – epidemiologically important
 - High-risk, high-volume procedures
 - Infections having known risk reduction methods



Targeted Surveillance

| Pros | Cons |
|---|--------------------------------------|
| Risk-adjusted rates | May miss some infections |
| Can measure trends and make comparisons | Limited information on endemic rates |
| More efficient use of resources | |
| Can target potential problems | |
| Identify performance improvement opportunities | |
| Can evaluate effectiveness of prevention activities | |

Combination

- Monitor:
 - Targeted events in defined populations and
 - Selected whole-house events
- Pros:
 - Rates are risk-adjusted
 - Measure trends
 - Target potential problems
 - Track selected events house-wide
- Cons:
 - May miss some infections



SPICE 

13


Selection of Processes and Outcomes

Processes-areas you might want to consider (CMS 8/24):

- Hand hygiene
- Appropriate use of PPE
- Point-of-care testing
- Urinary Catheter insertion/maintenance
- Cleaning and disinfection products/procedures

Outcomes

- Acute respiratory infections
- Urinary tract infections
- Skin/Soft Tissue Infections
- Gastroenteritis



SPICE 

14

Consideration for Choosing Outcome Measures

- Mandatory/required-Cat 1C
- *Frequency (incidence) of the infection
- *Communicability
- *System/resident cost (↑mortality, hospitalization)
- *Early Detection

**Based on the Infection Prevention risk assessment*



15

Should be included in routine surveillance

| Points to Consider | Infections | Comments |
|--|--|--|
| Evidence of <u>transmissibility</u> in a healthcare setting | Viral respiratory tract infections, viral GE, and viral conjunctivitis | Associated with outbreaks among residents and HCP in LTCFs |
| Processes available to prevent acquisition of infection, i.e., HH compliance | | |
| Clinically significant cause of morbidity or mortality | Pneumonia, UTI, GI tract infections, (including <i>C. difficile</i>) and SSTI | Associated with hospitalization and functional decline in LTCF residents |
| Specific pathogens causing serious outbreaks | Any invasive group A <i>Streptococcus</i> infection, acute viral hepatitis, norovirus, scabies, influenza-COVID-19, <i>C auris</i> | A single laboratory-confirmed case should prompt further investigation |



16

Infections that could be included in routine surveillance

| Points to Consider | Infections | Comments |
|---|--|---|
| Infections with limited transmissibility in a healthcare settings | Ear and sinus infections, fungal oral and skin infections and herpetic skin infections | Associated with underlying comorbid conditions and reactivation of endogenous infection |
| Infections with limited preventability | | |



17

Infections for which other accepted definitions should be applied in LTCF surveillance

| Points to Consider | Infections | Comments |
|---|---|--|
| Infections with other accepted definitions (may apply to only specific at-risk residents) | Surgical site infections, central-line- associated bloodstream infections and ventilator-associated pneumonia | LTCF-specific definitions were not developed. Refer to the National Healthcare Safety Network's criteria |



18

Sources of Data for Surveillance

- Clinical ward/unit rounds
- Medical Chart
- Lab reports
- Kardex/Patient Profile/Temperature logs
- Antibiotic Starts
- IT support



SPICE 

19

Surveillance

- ***The facility's surveillance system must include a data collection tool and the use of nationally-recognized surveillance criteria such as but not limited to, the CDC's National Healthcare Safety Network (NHSN) Long Term Care Criteria to define infections or revised McGeer criteria***

State Operations Manual
Appendix PP - Guidance to Surveyors for
Long Term Care Facilities

Table of Contents
(Rev. 08-2024)

SPICE 

20

Guiding Principles for Using Standardized Definitions

Infection surveillance only

Identify trends in a population

Applied retrospectively as it relates to clinical diagnosis/treatment

Limited elements

Focus on transmissible/preventable infections

Not for clinical decision making

21

Minimum Criteria for Initiation of Antibiotics in Long-Term Care Residents

Suspected Urinary Tract Infection

Loeb et al. Development of Minimum Criteria for the Initiation of Antibiotics in Residents of Long-Term Care Facilities: Results of a Consensus Conference. Inf Control Hosp Epi. 2001

22

www.cdc.gov/nhsn/ltc/uti/index.html

Centers for Disease Control and Prevention
CDC 24/7: Saving Lives. Protecting People™

Search The CDC

National Healthcare Safety Network (NHSN)

NHSN

NHSN Login

About NHSN

Enroll Here

Materials for Enrolled Facilities

Ambulatory Surgery Centers

Acute Care Hospitals/Facilities

Long-term Acute Care Hospitals/Facilities

Long-term Care Facilities

Surveillance for C. difficile and MRSA Infections

Surveillance for C. difficile, MRSA, and other Drug-resistant Infections

Resources for NHSN Users Already Enrolled

Training

Protocol

Data Collection Forms

Supporting Material

Analysis Resources

New Users - Start Here

Step 1: Enroll into NHSN

Step 2: Set up NHSN

Step 3: Report

[Click here to enroll!](#)

5:22 PM 9/13/2016

23

Purposes of NHSN

- Provide facilities with risk-adjusted data that can be used for inter-facility comparisons and local quality improvement activities
- Assist facilities in developing surveillance and analysis methods that permit timely recognition of patient and healthcare personnel safety problems and prompt intervention with appropriate measures
- Conduct collaborative research studies with members
- Data repository for reporting healthcare associated infections, COVID-19 infections, COVID vaccinations and certain process measures

SPICE

24

Surveillance Definitions of Infections in Long-Term Care Facilities: Revisiting the McGeer Criteria

Nimalie D. Stone, MD;¹ Muhammad S. Ashraf, MD;² Jennifer Calder, PhD;³ Christopher J. Crnich, MD;⁴
 Kent Crossley, MD;⁵ Paul J. Drinka, MD;⁶ Carolyn V. Gould, MD;⁷ Manisha Juthani-Mehta, MD;⁷
 Ebbing Lautenbach, MD;⁸ Mark Loeb, MD;⁹ Taranisia MacCannell, PhD;¹ Preeti N. Malani, MD;^{10,11} Lona Mody, MD;^{10,11}
 Joseph M. Mylotte, MD;¹² Lindsay E. Nicolle, MD;¹³ Mary-Claire Roghmann, MD;¹⁴ Steven J. Schweon, MSN;¹⁵
 Andrew E. Simor, MD;¹⁶ Philip W. Smith, MD;¹⁷ Kurt B. Stevenson, MD;¹⁸ Suzanne F. Bradley, MD^{10,11}
 for the Society for Healthcare Epidemiology Long-Term Care Special Interest Group*



Long-term Care Catheter-associated UTI
 Infection Worksheet
 8/6/09; Criteria-2012

| Resident Name | MR # | Date of Birth | Resident Location (Hall/room#) |
|---|------|--|---|
| Date Specimen collected: | | Gender: | Resident type: <input type="checkbox"/> Short-stay <input type="checkbox"/> Long-stay |
| <input type="checkbox"/> < 2 calendar days Community-Onset (CO) | | <input type="checkbox"/> Male | Date of 1 st admission to facility: / / |
| <input type="checkbox"/> > 2 calendar days Long-term Care Facility Onset (LO) | | <input type="checkbox"/> Female | Date of current admission to facility: / / |
| Primary Resident Service Types: | | Has resident been transferred from an acute care facility to your facility in the past 4 weeks? <input type="checkbox"/> Yes <input type="checkbox"/> No | |
| <input type="checkbox"/> Long-term general nursing | | *If yes, date of last transfer from acute care to your facility: / / | |
| <input type="checkbox"/> Long-term dementia <input type="checkbox"/> Long-term psychiatric | | *If yes, did the resident have an indwelling catheter at the time of transfer to your facility? <input type="checkbox"/> Yes <input type="checkbox"/> No | |
| <input type="checkbox"/> Skilled nursing/Short-term Rehab <input type="checkbox"/> Bariatric | | If urinary catheter in place or removed within last 2 calendar days: | |
| <input type="checkbox"/> Hospice/Palliative <input type="checkbox"/> Ventilator | | Site where inserted: <input type="checkbox"/> Your facility <input type="checkbox"/> Hospital <input type="checkbox"/> Other | |
| Indwelling Urinary Catheter status at time of event onset: | | Date of urinary catheter insertion: / / | |
| <input type="checkbox"/> In place <input type="checkbox"/> Removed within last 2 calendar days | | Transfer to acute care facility within 7 days? <input type="checkbox"/> Yes <input type="checkbox"/> No | |
| <input type="checkbox"/> Not in place | | If urinary catheter not in place, was there another urinary device type present at time of event onset? <input type="checkbox"/> Yes <input type="checkbox"/> No | |
| If Yes, other device type: <input type="checkbox"/> Suprapubic <input type="checkbox"/> Condom (males only) <input type="checkbox"/> Intermittent straight catheter | | Person completing form: / / | |
| Date of Event (date of first sign/symptom OR date of specimen): / / | | | |

Criteria for Symptomatic Urinary Tract Infection, with an Indwelling catheter (Ca-UTI)

For residents with an indwelling catheter both Criteria 1 and 2 must be met

Criteria 1
 At least one of the following sign/symptom sub-criteria presents:
 a) Fever, rigors, or new-onset hypotension with no alternate site of infection
 b) Either acute change in mental status or functional decline, with no alternate diagnosis AND leukocytosis
 c) New-onset suprapubic pain or costovertebral angle pain or tenderness
 d) Purulent discharge from around the catheter or acute pain, swelling, or tenderness of the testes, epididymis, or prostate

Criteria 2
 Urinary catheter specimen culture with at least:
 At least 10⁵ cfu/ml (>100,000 cfu) of any organism(s)

Surveillance Definitions of Infections in Long-Term Care Facilities: Revisiting the McGeer Criteria
<http://www.iche.ohsu.edu/guidelines/inf/inf120809/inf12080901010.pdf>

Long-term Care Catheter-associated UTI
 Infection Worksheet
 8/6/09; Criteria-2012

Comments:

1. Urine specimens should be processed within 1-2 hours OR refrigerated and processed within 24 hours.
2. Recent catheter trauma, catheter obstruction or new onset of hematuria are useful localizing signs that are consistent with UTI but are not necessary for diagnosis
3. Urinary catheter specimens for culture should be collected following replacement of the catheter (if current catheter has been in place for >14 days)

Surveillance Definitions of Infections in Long-Term Care Facilities: Revisiting the McGeer Criteria
<http://www.iche.ohsu.edu/guidelines/inf/inf120809/inf12080901010.pdf>



Attribution of infection to LTCF

- No evidence of an incubating infection at the time of admission to the facility
 - Basis of clinical documentation of appropriate signs and symptoms and not solely on screening microbiologic data
- Onset of clinical manifestation occurs > 2 calendar days after admission.



SPICE 

27

Attribution of infection to LTCF

- All symptoms must be new or acutely worse
- Non-infectious causes of signs and symptoms should always be considered prior to diagnosis
- Identification of an infection should not be based on a single piece of evidence
 - Clinical, microbiologic, radiologic
- Diagnosis by physician insufficient (based on definition)


SPICE 

28

Constitutional Requirements

Fever:

- A single oral temperature $>37.8^{\circ}\text{C}$ [100°F], OR
- Repeated oral temperatures $>37.2^{\circ}\text{C}$ [99°F]; rectal temperature $>37.5^{\circ}$ (99.5°F) OR
- $>1.1^{\circ}\text{C}$ [2°F] over baseline from a temperature taken at any site

SPICE 

29

Constitutional Requirements

Leukocytosis

- Neutrophilia > 14000 WBC/ mm^3
- OR
- Left shift ($>6\%$ bands or ≥ 1500 bands/ mm^3)

SPICE 

30

Constitutional Requirements

Acute Change in Mental Status from Baseline

- Based on Confusion Assessment Method (CAM) criteria available in MDS

| Change | Criteria |
|--------------------------------|---|
| Acute Onset | Evidence of acute change in mental status from resident baseline |
| Fluctuating | Behavior fluctuating (e.g., coming and going or changing in severity during assessment) |
| Inattention | Resident has difficulty focusing attention (e.g., unable to keep track of discussion or easily distracted) |
| Disorganized Thinking | Resident's thinking is incoherent (e.g., rambling conversation, unclear flow of ideas) |
| Altered level of consciousness | Resident's level of consciousness is described as different from baseline (e.g., hyperalert, sleepy, drowsy, difficult arouse, nonresponsive) |

Either /or



31

Constitutional Requirements

Acute Functional Decline

- New 3-point increase in total ADL score (0-28) from baseline based on 7 ADLs {0 = independent; 4 = total dependence}
 - Bed mobility
 - Transfer
 - Locomotion within LTCF
 - Dressing
 - Toilet use
 - Personal hygiene
 - Eating



32

Site Specific Definitions

Knowledge Checks





SPICE 

33

Respiratory Tract Infections

| Criteria | Comments |
|---|---|
| <p>A. <i>Common cold syndrome/pharyngitis</i></p> <p>At least two criteria present</p> <ol style="list-style-type: none"> 1. Runny nose or sneezing 2. Stuffy nose (i.e., congestion) 3. Sore throat or hoarseness or difficulty swallowing 4. Dry cough 5. Swollen or tender glands in neck | <p>Fever may or may not be present. Symptoms must be new, and not attributable to allergies</p> |

SPICE 

34

Respiratory Tract Infections

| Criteria | Comments |
|--|--|
| <p>B. Influenza-like Illness</p> <p>Both criteria 1 and 2 present</p> <ol style="list-style-type: none"> 1. Fever 2. At least three of the following symptom sub-criteria (a-f) present <ol style="list-style-type: none"> a. Chills b. New headache or eye pain c. Myalgias or body aches d. Malaise or loss of appetite e. Sore throat f. New or increased dry cough | <p>If criteria for influenza-like illness and another upper or lower respiratory tract infection are met at the same time, only the diagnosis of influenza-like illness should be used</p> <p>Due to increasing uncertainty surrounding the timing of the start of influenza season, the peak of influenza activity and the length of the season, 'seasonality' is no longer part of the criteria to define influenza-like illness</p> |

35


Respiratory Tract Infections

| Criteria | Comments |
|--|---|
| <p>C. Pneumonia</p> <p>All criteria 1-3 present</p> <ol style="list-style-type: none"> 1. Interpretation of chest radiograph as demonstrating pneumonia or the presence of <u>new</u> infiltrate 2. At least one of the following respiratory sub-criteria (a-f) present <ol style="list-style-type: none"> a. New or increased cough b. New or increased sputum production c. O₂ saturation <94% on room air or a reduction in O₂ saturation of more than 3% from baseline d. New or changed lung exam abnormalities e. Pleuritic chest pain f. Respiratory rate of \geq 25/min 3. At least one constitutional criteria | <p>For both pneumonia and lower respiratory tract infections, presence of underlying conditions which could mimic a respiratory tract infection presentation (congestive heart failure, interstitial lung disease), should be excluded by review of clinical records and an assessment of presenting symptoms and signs</p> |



36

Respiratory Tract Infections


| Criteria | Comments |
|---|---|
| <p>D. Lower respiratory tract (Bronchitis or Tracheo-bronchitis)</p> <p>All criteria 1-3 present</p> <ol style="list-style-type: none"> 1. Chest radiograph not performed <u>or</u> <i>negative</i> for pneumonia or new infiltrate. 2. At least two of the following respiratory sub-criteria (a-f) present <ol style="list-style-type: none"> a. New or increased cough b. New or increased sputum production c. O₂ saturation <94% on room air or a reduction in O₂ saturation of more than 3% from baseline d. New or changed lung exam abnormalities e. Pleuritic chest pain f. Respiratory rate of ≥ 25/min 3. At least one constitutional criteria | <p>For both pneumonia and lower respiratory tract infections, presence of underlying conditions which could mimic a respiratory tract infection presentation (congestive heart failure, interstitial lung disease), should be excluded by review of clinical records and an assessment of presenting symptoms and signs</p> |



37

Knowledge Check




38

Knowledge Check #1

Mr. Do Little has multiple co-morbidities including hypertension and acute respiratory failure. Vitals on admission WNL

On day seven after admission, the daughter tells the nurse “dad is not responding like he used to. He can not hold a conversation, tires easily and is not able to brush his teeth, eat or dress without assistance.” **I think he has a UTI. He needs an antibiotic.**



39

Clinical Picture


Physical exam:

- Temp 100.7, pulse 107, RR 26 and O2 sat 93%
- Ronchi noted on auscultation of the chest the resident is confused

MD notified and orders urine and chest x-ray

Results:

- Culture + E. coli 10² cfu/ml and
- chest x-ray: no new findings



40

Does Mr. Do Little have an infection?

- Yes
- No
- Have no idea

SPICE 

41

What surveillance criteria are met?

- A. Common Cold
- B. Pneumonia
- C. Urinary tract infection
- D. Lower respiratory track

SPICE 

42

If Yes, is it facility or community associated?

- Facility
- Community




43

Respiratory Tract Infections

| Criteria | Comments |
|---|---|
| <p>D. <u>Lower respiratory tract (Bronchitis or Tracheo-bronchitis)</u></p> <p>All criteria 1-3 present</p> <ol style="list-style-type: none"> 1. Chest radiograph not performed <u>or negative for pneumonia or new infiltrate.</u> 2. At least two of the following respiratory sub-criteria (a-f) present <ol style="list-style-type: none"> a. New or increased cough b. New or increased sputum production c. <u>O₂ saturation <94% on room air or a reduction in O₂ saturation of more than 3% from baseline</u> d. <u>New or changed lung exam abnormalities</u> e. Pleuritic chest pain f. Respiratory rate of ≥ 25/min 3. At least one constitutional criteria | <p>For both pneumonia and lower respiratory tract infections, presence of underlying conditions which could mimic a respiratory tract infection presentation (congestive heart failure, interstitial lung disease), should be excluded by review of clinical records and an assessment of presenting symptoms and signs</p> |

44

WHAT ABOUT URINARY TRACT INFECTIONS



45

What do the Guidelines Say?

- Insert catheters only for appropriate indications
- Avoid use of urinary catheters in patients and nursing home residents for management of incontinence
- Keep the catheter and collecting tube free from kinking
- Empty the drainage bag regularly using a separate, clean collecting container for each resident (even in semi-private rooms)
- **Changing indwelling catheters or drainage bags at routine, fixed intervals is not recommended.** It is suggested to change catheters and drainage bags based on clinical indications such as infection, obstruction, or when the closed system has been compromised

<https://www.cdc.gov/infectioncontrol/guidelines/cauti/index.html>




46

Urinary Specimens: What do the Guidelines Say?

- Specimens collected through the catheter present for more than a few days reflect biofilm microbiology.
- For residents with chronic indwelling catheters (greater than 14 days) and symptomatic infection, changing the catheter immediately prior to instituting antimicrobial therapy allows collection of a bladder specimen, which is a more accurate reflection of infecting organisms.
- Urinary catheters coated with antimicrobial materials have the potential to decrease UTIs but have not been studied in the LTCF setting.

SHEA/APIC Guideline: Infection prevention and control in the long-term care facility Philip W. Smith, MD, Gail Bennett, RN, MSN, CICb Suzanne Bradley, MD, Paul Drinka, MD, Ebbing Lautenbach, MD, James Marx, RN, MS, CIC, Lona Mody, MD, Lindsay Nicolle, MD and Kurt Stevenson, MD July 2008



47

| McGeer Urinary Tract Infections <i>without catheter</i> | | |
|---|--|--|
| Criteria | | Comments |
| <p>A. For Residents without an indwelling catheter</p> <p>Both criteria 1 and 2 present</p> <p>1. At least one of the following sign/symptom sub-criteria (a-c) present:</p> <div style="border: 1px solid black; padding: 5px; margin-bottom: 5px;"> <p>a) Acute dysuria <u>or</u> acute pain, swelling, or tenderness of the testes, epididymis, or prostate</p> </div> <div style="border: 1px solid black; padding: 5px;"> <p>b) Fever <u>or</u> leukocytosis and</p> <p>At least one of the following localizing urinary tract sub-criteria:</p> <ul style="list-style-type: none"> i. Acute costovertebral angle pain or tenderness ii. Suprapubic pain iii. Gross hematuria iv. New or marked increase in incontinence v. New or marked increase in urgency vi. New or marked increase in frequency </div> | <p>c) In the absence of fever of leukocytosis, then at least two or more of the following localizing urinary symptoms</p> <ul style="list-style-type: none"> i. Suprapubic pain ii. Gross hematuria iii. New or marked increase in incontinence iv. New or marked increase in urgency v. New or marked increase in frequency <p>2. One of the following microbiologic sub-criteria</p> <div style="border: 1px solid black; padding: 5px; margin-bottom: 5px;"> <p>a) At least 10⁵ cfu/ml of no more than 2 species of microorganisms in a voided urine</p> </div> <div style="border: 1px solid black; padding: 5px;"> <p>b) At least 10² cfu/ml of any number of organisms in a specimen collected by an in and out catheter</p> </div> | <p>UTI should be diagnosed when there are localizing s/s <u>and</u> a positive urinary culture</p> <p><i>A diagnosis of UTI can be made without localizing symptoms if a blood culture isolate of the same organism isolated from the urine and there is no alternate sight of infection</i></p> <p>In the absence of a clear alternate source, fever or rigors with a positive urine culture in a non-catheterized resident will often be treated as a UTI. However, evidence suggest most of these episodes are not from a urinary source</p> <p>Pyuria does not differentiate symptomatic UTI from asymptomatic bacteria</p> <p>Absence of pyuria in diagnostic test excludes symptomatic UTI in residents of LTCF</p> <p>Urine specimens should be processed within 1-2 hours, or refrigerated and processed with in 24 hours.</p> |

48

McGeer Urinary Tract Infections

| Criteria | Comments |
|--|---|
| <p>B. For the resident with an indwelling catheter</p> <p>Both criteria 1 and 2 present</p> <p>1. At least one of the following sign/symptom sub-criteria (a-d) present:</p> <ul style="list-style-type: none"> a) Fever, rigors, or new onset hypotension, with no alternate site of infection b) Either acute change in mental status <u>or</u> acute functional decline with no alternate diagnosis <u>and</u> Leukocytosis c) New onset suprapubic pain <u>or</u> costovertebral angle pain or tenderness d) Purulent discharge from around the catheter <u>or</u> acute pain, swelling, or tenderness of the testes, epididymis, or prostate <p>2. Urinary catheter culture with at least 10⁵ cfu/ml of any organism(s)</p> | <p>Recent catheter trauma, catheter obstruction or new onset hematuria are useful localizing signs consistent with UTI, but not necessary for diagnosis</p> <p>Urinary catheter specimens for culture should be collected following the replacement of the catheter (if current catheter has been in place for >14 days)</p> |

49



NHSN

National Healthcare Safety Network



50

NHSN Notes


- Indwelling urinary catheter should be in place for a minimum of 2 calendar days before infection onset (day 1 = day of insertion)
- Indwelling urinary catheter: a drainage tube that is inserted into the urinary bladder through the urethra, is left in place and is connected to a closed collection system, also called a foley catheter. Indwelling urinary catheters do not include straight in-and-out catheters or suprapubic catheters (these would be captures as SUTIs, not CA-SUTIs)
- *Indwelling catheters which have been in place for > 14 days should be changed prior to specimen collection but failure to change catheter does not exclude a UTI for surveillance purposes*



51

NHSN Key Reminders

1. ***“Mixed flora” is not available in the pathogen list within NHSN.*** Therefore, it cannot be reported as a pathogen to meet the NHSN UTI criteria. Additionally, “mixed flora” often represents contamination and likely represents presence of multiple organisms in culture (specifically, at least two organisms).
2. ***Yeast and other microorganisms, which are not bacteria, are not acceptable UTI pathogens,*** and therefore, cannot be used to meet NHSN UTI criteria without the presence of a qualifying bacterium.
3. To remove the subjectivity about whether a fever is attributable to a UTI event, ***the presence of a fever, even if due to another cause (for example, pneumonia), must still be counted as a criterion when determining if the NHSN UTI definition is met.***



52

| | | | |
|--|---|--|--|
| <h2 style="margin: 0;">NHSN Urinary Tract Infections</h2> <h3 style="margin: 0;">For Residents without an indwelling catheter</h3> | Must meet Criteria 1,2 OR 3 | | |
| | <p>Criteria 1</p> <p>One of the following true:</p> <ol style="list-style-type: none"> 1. Acute dysuria 2. Acute pain, swelling or tenderness of the testes, epididymis or prostate <p style="text-align: center;">AND</p> <p>A positive urine culture with no more than 2 species of microorganisms, at least one of which is a bacterium of $\geq 10^5$ CFU/ml</p> | <p>Criteria 2</p> <p>Either of the following:</p> <ol style="list-style-type: none"> 1. Fever: (Single temperature $>100^\circ$ F or $>99^\circ$ F on repeated occasions (more than once) OR an increase of $>2^\circ$ F over baseline 2. Leukocytosis: $>10,000$ cells/mm³ or left shift (6% or 1,500 bands/mm³) <p style="text-align: center;">AND</p> <p>One or more of the following (New or Marked increase):</p> <ol style="list-style-type: none"> 1. Costovertebral angle pain or tenderness 2. Suprapubic tenderness 3. Visible (Gross) hematuria 4. Incontinence 5. Urgency 6. Frequency <p style="text-align: center;">AND</p> <p>A positive urine culture with no more than 2 species of microorganisms, at least one of which is a bacterium of $\geq 10^5$ CFU/m</p> | <p>Criteria 3</p> <p>Two or more of the following (New and/or marked increase):</p> <ol style="list-style-type: none"> 1. Costovertebral angle pain or tenderness 2. Incontinence 3. Urinary urgency 4. Urinary frequency 5. Suprapubic tenderness 6. Visible (gross) hematuria <p style="text-align: center;">AND</p> <p>A positive urine culture with no more than 2 species of microorganisms, at least one of which is a bacterium of $\geq 10^5$ CFU/ml</p> |
| <p><i>Comments: Fever can be used to meet SUTI criteria even if the resident has another possible cause for the fever (for example, pneumonia)</i></p> | | | |

53

| <h2 style="margin: 0;">NHSN Urinary Tract Infections</h2> | |
|---|---|
| <p style="font-size: x-small; margin: 0;">For the resident with an indwelling catheter-or removed within 2 calendar days prior to the event onset (day of catheter removal is day 1)</p> | |
| Criteria | Comments |
| <p>CA-SUTI</p> <p>One or more of the following (Signs and Symptoms and Laboratory and diagnostic Testing):</p> <ol style="list-style-type: none"> a) *Fever (Single temperature $>100^\circ$ F or $>99^\circ$ F on repeated occasions OR an increase of $>2^\circ$ F over baseline) b) Rigors c) New onset hypotension, with no alternate noninfectious cause d) New onset confusion/functional decline with no alternate diagnosis AND Leukocytosis ($>10,000$ cells/mm³ or left shift (6% or 1,500 bands/mm³)) e) New or marked increase in costovertebral angle pain or tenderness f) New or marked increase in suprapubic tenderness g) Acute pain, swelling, or tenderness of the testes, epididymis, or prostate h) Purulent discharge from around the catheter insertion site i) Acute dysuria (used only if catheter removed) | <p style="font-size: 2em; font-weight: bold; margin: 0;">AND</p> <div style="text-align: center;"> </div> <p style="color: red; font-weight: bold; margin: 0;">A positive urine culture with no more than 2 species of microorganisms, at least one of which is a bacterium of $\geq 10^5$ CFU/ml</p> <p style="font-size: small; margin-top: 20px;"><i>*Fever can be used to meet CA-SUTI criteria even if the resident has another possible cause for the fever (for example, pneumonia)</i></p> |

54

Asymptomatic Bacteremic Urinary Tract Infection (ABUTI)

Resident with or without an indwelling urinary catheter:

1. No qualifying fever or signs or symptoms (specifically no urinary urgency, urinary frequency, acute dysuria, suprapubic tenderness, or costovertebral angle pain or tenderness). If no catheter is in place, fever alone would not exclude ABUTI if other criteria are met

AND

2. A positive urine culture with no more than 2-species of microorganisms, at least one of which is a bacterium of $\geq 10^5$ CFU/ml


AND

3. A positive blood culture with at least 1 matching bacteria to the urine culture

SPICE 

55

Knowledge Check



SPICE 

56

Knowledge Check # 1



1 Mar.

Mrs. Ross is a resident in your facility, admitted on February 1st. An indwelling urinary catheter was inserted on March 1st.



5 Mar.

On March 5, the nurse practitioner documented that Mrs. Ross complained of suprapubic pain.



6 Mar.

The following day, on March 6, a specimen collected from the Foley catheter was sent to the lab and subsequently tested positive for greater than 100,000 CFU/ml of E. coli and 100,000 CFU/ml of Candida



57

Does Mr. Ross have a UTI?

A. Yes

B. No



58

If Yes, is it catheter associate?

A. No

B. Yes

What criteria are met?



59

McGeer Urinary Tract Infections

| Criteria | Comments |
|---|---|
| B. For the resident with an indwelling catheter | |
| Both criteria 1 and 2 present | Recent catheter trauma, catheter obstruction or new onset hematuria are useful localizing signs consistent with UTI, but not necessary for diagnosis |
| 1. At least one of the following sign/symptom sub-criteria (a-d) present: | |
| a) Fever, rigors, or new onset hypotension, with no alternate site of infection | |
| b) Either acute change in mental status <u>or</u> acute functional decline with no alternate diagnosis <u>and</u> Leukocytosis | |
| c) New onset suprapubic pain <u>or</u> costovertebral angle pain or tenderness | ✓ Urinary catheter specimens for culture should be collected following the replacement of the catheter (if current catheter has been in place for >14 days) |
| d) Purulent discharge from around the catheter <u>or</u> acute pain, swelling, or tenderness of the testes, epididymis, or prostate | |
| 2. Urinary catheter culture with $\geq 10^5$ cfu/ml of any organism(s) | ✓ |



60

| NHSN Urinary Tract Infections | |
|---|---|
| <i>For the resident with an indwelling catheter</i> ✓ | |
| Criteria | Comments |
| <u>CA-SUTI</u> | |
| <p>One or more of the following (Signs and Symptoms and Laboratory and diagnostic Testing):</p> <ul style="list-style-type: none"> a) *Fever (Single temperature >100° F or >99° F on repeated occasions OR an increase of >2° F over baseline) b) Rigors c) New onset hypotension, with no alternate site of infection d) New onset confusion/functional decline AND Leukocytosis (>14,000 cells/mm³ or left shift (6% or 1,500 bands/mm³)) e) New or marked increase in costovertebral angle pain or tenderness f) New or marked increase in suprapubic tenderness ✓ g) Acute pain, swelling, or tenderness of the testes, epididymis, or prostate h) Purulent discharge from around the catheter | <p>AND</p> <p>↓</p> <p>A positive urine culture with no more than 2 species of microorganisms, at least one of which is a bacterium of $\geq 10^5$ CFU/ml ✓</p> <p><i>*Fever can be used to meet CA-SUTI criteria even if the resident has another possible cause for the fever (for example, pneumonia)</i></p> |

61


Knowledge Check # 2

Mr. U, a resident of LTC facility has a urinary catheter in place for 3 days for acute urinary retention. Later that day he spikes a fever of 101°F and has a cough with shortness of breath.

The physician orders a urine culture, and it comes back positive with >100,000 CFU/ml of Pseudomonas aeruginosa and Candida albicans.

Upon further work, up Mr. U is determined not to have any other symptoms that meet the NHSN CA-SUTI criteria,

§ But, a chest X-ray does show infiltrates in the right upper lobe of the lung.



62

Does Mr. U have an infection?

- A. Yes
- B. No

What type of infection does Mr. U have?

- 1. Pneumonia
- 2. Catheter Associated UTI
- 3. UTI but no catheter



63

McGeer Urinary Tract Infections

| Criteria | Comments |
|---|---|
| <p>B. For the resident with an indwelling catheter</p> <p>Both criteria 1 and 2 present</p> <p>1. At least one of the following sign/symptom sub-criteria (a-d) present:</p> <ul style="list-style-type: none"> a) Fever, rigors, or new onset hypotension, with no alternate site of infection ✓ b) Either acute change in mental status <u>or</u> acute functional decline with no alternate diagnosis <u>and</u> Leukocytosis c) New onset suprapubic pain <u>or</u> costovertebral angle pain or tenderness d) Purulent discharge from around the catheter <u>or</u> acute pain, swelling, or tenderness of the testes, epididymis, or prostate <p>2. Urinary catheter culture with $\geq 10^5$ cfu/ml of any organism(s) ✓</p> | <p>Recent catheter trauma, catheter obstruction or new onset hematuria are useful localizing signs consistent with UTI, but not necessary for diagnosis</p> <p>Urinary catheter specimens for culture should be collected following the replacement of the catheter (if current catheter has been in place for >14 days)</p> |



64

NHSN Urinary Tract Infections

*For the resident with an **indwelling catheter** ✓ > 2 days*

| Criteria | Comments |
|--|---|
| <i>CA-SUTI</i> | |
| One or more of the following (Signs and Symptoms and Laboratory and diagnostic Testing): ✓ | <p>AND</p> <p>↓</p> <p>A positive urine culture with no more than 2 species of microorganisms, at least one of which is a bacterium of $\geq 10^5$ CFU/ml ✓</p> <p><i>*Fever can be used to meet CA-SUTI criteria even if the resident has another possible cause for the fever (for example, pneumonia)</i></p> |
| a) *Fever (Single temperature >100° F or >99° F on repeated occasions OR an increase of >2° F over baseline) ✓ | |
| b) Rigors | |
| c) New onset hypotension, with no alternate site of infection | |
| d) New onset confusion/functional decline AND Leukocytosis (>14,000 cells/mm ³ or left shift (6% or 1,500 bands/mm ³)) | |
| e) New or marked increase in costovertebral angle pain or tenderness | |
| f) New or marked increase in suprapubic tenderness | |
| g) Acute pain, swelling, or tenderness of the testes, epididymis, or prostate | |
| h) Purulent discharge from around the catheter | |

65

Respiratory Tract Infections

| Criteria | Comments |
|---|---|
| C. Pneumonia | |
| All criteria 1-3 present | |
| 1. Interpretation of chest radiograph as demonstrating pneumonia or the presence of new infiltrate ✓ | <p>For both pneumonia and lower respiratory tract infections, presence of underlying conditions which could mimic a respiratory tract infection presentation (congestive heart failure, interstitial lung disease), should be excluded by review of clinical records and an assessment of presenting symptoms and signs</p> |
| 2. At least one of the following respiratory sub-criteria (a-f) present | |
| a. New or increased cough ✓ | |
| b. New or increased sputum production | |
| c. O ₂ saturation <94% on room air or a reduction in O ₂ saturation of more than 3% from baseline | |
| d. New or changed lung exam abnormalities | |
| e. Pleuritic chest pain | |
| f. Respiratory rate of \geq 25/min | |
| 3. At least one constitutional criteria ✓ | |



66

Knowledge Check # 3

Day 1: Ms. R had an indwelling urinary catheter inserted for a bladder outlet obstruction

Day 2: The indwelling urinary catheter remains in place

Day 3: The resident's indwelling urinary catheter remains in place. The resident had a single oral temp of 100.2°F. A urine culture was collected from the catheter



67

Ms. R
continued

Day 4: The indwelling urinary catheter remains in place. No symptoms documented

Day 5: The urine culture was positive for *Candida glabrata* 10^5 CFU/ml



68

Does Ms. R have an infection?

A. Yes

B. No

What type of infection does Ms. R have?

1. Catheter Associated UTI
2. UTI but no catheter

Which definition is met?

1. NHSN
2. McGeer
3. Both



69

McGeer Urinary Tract Infections

| Criteria | Comments |
|---|---|
| <p>B. For the resident with an indwelling catheter ✓</p> <p>Both criteria 1 and 2 present</p> <p>1. At least one of the following sign/symptom sub-criteria (a-d) present:</p> <ol style="list-style-type: none"> a) Fever, rigors, or new onset hypotension, with no alternate site of infection ✓ b) Either acute change in mental status <u>or</u> acute functional decline with no alternate diagnosis <u>and</u> Leukocytosis c) New onset suprapubic pain <u>or</u> costovertebral angle pain or tenderness d) Purulent discharge from around the catheter <u>or</u> acute pain, swelling, or tenderness of the testes, epididymis, or prostate <p>2. Urinary catheter culture with $\geq 10^5$ cfu/ml of any organism(s) ✓</p> | <p>Recent catheter trauma, catheter obstruction or new onset hematuria are useful localizing signs consistent with UTI, but not necessary for diagnosis</p> <p>Urinary catheter specimens for culture should be collected following the replacement of the catheter (if current catheter has been in place for >14 days)</p> |



70

Knowledge Check # 4

Mrs. C is an 85-year-old female who is normally ambulatory, independent of ADLs and very social with staff and other residents. She has been a resident of your facility for 10 years

This morning, March 5th, Mrs. C seems confused, refuses breakfast, is incontinent of stool and does not want to get out of bed.

Vital Signs: Temp 99.5, RR 22, O²Sat 93% on room air and BP is 110/70. Urine is dark yellow and has a strong odor.

Physician orders, UC, BC and chest x-ray

SPICE 

71

Knowledge Check # 4

Diagnostic test are completed, and results are as follows:

UC positive for $>10^5$ cfu/ml of *klebsiella pneumonia* and $> 10^2$ *candida albicans*

Chest x-ray negative for infiltrate

BC + for *Klebsiella pneumonia*

What Surveillance Definition Does Mrs. C meet?

1. Lower respiratory tract
2. Gastroenteritis
3. Urinary tract infection
4. Bloodstream infection
5. Asymptomatic Bacteremic Urinary Tract Infection

SPICE 

72

Asymptomatic Bacteremic Urinary Tract Infection (ABUTI)

Resident with or without an indwelling urinary catheter:

1. **No qualifying fever or signs or symptoms** (specifically no urinary urgency, urinary frequency, acute dysuria, suprapubic tenderness, or costovertebral angle pain or tenderness). If no catheter is in place, fever alone would not exclude ABUTI if other criteria are met
- AND**
2. A **positive urine culture** with no more than 2-species of microorganisms, at least one of which is a bacterium of $\geq 10^5$ CFU/ml
- AND**
3. A **positive blood culture with at least 1 matching bacteria** to the urine culture

SPICE 

73

Skin, Soft Tissue and Mucosal Infections



SPICE 

74

Skin, Soft Tissue and Mucosal Infections

| Criteria | Comments |
|--|---|
| <p>A. <u>Cellulitis/soft tissue/wound infection</u></p> <p>At least one of the following criteria is present</p> <ol style="list-style-type: none"> 1. Pus present at a wound, skin, or soft tissue site 2. New or increasing presence of at least four of the following sign/symptom sub-criteria <ol style="list-style-type: none"> a) Heat at affected site b) Redness at affected site c) Swelling at affected site d) Tenderness or pain at affected site e) Serous drainage at affected site f) One constitutional criteria | <p>More than one resident with streptococcal skin infection from the same serogroup (e.g., A, B, C, G) in a LTCF may suggest an outbreak</p> <p>For wound infections related to surgical procedures: LTCF should use the CDC's NHSN surgical site infection criteria and report these infections back to the institution performing the original surgery</p> <p>Presence of organisms cultured from the surface (e.g., superficial swab culture) of a wound is not enough evidence that the wound is infected</p> |

75

Skin, Soft Tissue and Mucosal Infections

| Criteria | Comments |
|---|--|
| <p>B. <u>Scabies</u></p> <p>Both criteria 1 and 2 present</p> <ol style="list-style-type: none"> 1. A maculopapular and/or itching rash 2. At least one of the following sub-criteria: <ol style="list-style-type: none"> a) Physician diagnosis b) Laboratory confirmation (scrapping or biopsy) c) Epidemiologic linkage to a case of scabies with laboratory confirmation | <p>Care must be taken to rule out rashes due to skin irritation, allergic reactions, eczema, and other non-infectious skin conditions</p> <p>An epidemiologic linkage to a case can be considered if there is evidence of geographic proximity in the facility, temporal relationship to the onset of symptoms, or evidence of a common source of exposure (i.e., shared caregiver).</p> |

76

Skin, Soft Tissue and Mucosal Infections

| Criteria | Comments |
|---|--|
| <p>C. <u>Fungal oral/perioral and skin infections</u></p> <p><u>Oral candidiasis:</u> Both criteria 1 and 2 present:</p> <ol style="list-style-type: none"> 1. Presence of raised white patches on inflamed mucosa, or plaques on oral mucosa 2. Medical or dental provider diagnosis <p><u>Fungal skin Infection:</u> Both criteria 1 and 2 present:</p> <ol style="list-style-type: none"> 1. Characteristic rash or lesion 2. Either a medical provider diagnosis or laboratory confirmed fungal pathogen from scraping or biopsy | <p>Mucocutaneous candida infections are usually due to underlying clinical conditions such as poorly controlled diabetes or severe immunosuppression. Although not transmissible infections in the healthcare setting, they can be a marker for increased antibiotic exposure</p> <p>Dermatophytes have been known to cause occasional infections, and rare outbreaks, in the LTC setting.</p> |

77

Skin, Soft Tissue and Mucosal Infections

| Criteria | Comments |
|---|--|
| <p>D. <u>Herpes viral skin infections</u></p> <p><u>Herpes simplex infection</u> Both criteria 1 and 2 present:</p> <ol style="list-style-type: none"> 1. A vesicular rash 2. Either physician diagnosis or laboratory confirmation <p><u>Herpes zoster infection</u> Both criteria 1 and 2 present:</p> <ol style="list-style-type: none"> 1. A vesicular rash 2. Either physician diagnosis or laboratory confirmation | <p>Reactivation of old herpes simplex ("cold sores") or herpes zoster ("shingles") is not considered a healthcare-associated infection</p> <p>Primary herpes viral skin infections are very uncommon in LTCF, except in pediatric populations where it should be considered healthcare-associated.</p> |

78


Skin, Soft Tissue and Mucosal Infections

| Criteria | Comments |
|---|---|
| <p>E. <i>Conjunctivitis</i></p> <p>At least one of the following criteria present:</p> <ol style="list-style-type: none"> 1. Pus appearing from one or both eyes, present for at least 24 hours 2. New or increasing conjunctival erythema, with or without itching. 3. New or increased conjunctival pain, present for at least 24 hours. | <p>Conjunctivitis symptoms (“pink eye”) should not be due to allergic reaction or trauma.</p> |



79

Gastrointestinal Tract Infections



80

Gastrointestinal Tract Infections

| Criteria | Comments |
|--|---|
| <p>A. <u>Gastroenteritis</u></p> <p>At least one of the following criteria present</p> <ol style="list-style-type: none"> <li style="margin-bottom: 5px;">1. Diarrhea, three or more liquid or watery stools above what is normal for the resident within a 24-hour period <li style="margin-bottom: 5px;">2. Vomiting, two or more episodes in a 24-hour period <li style="margin-bottom: 5px;">3. Both of the following sign/symptom sub-criteria present: <ol style="list-style-type: none"> <li style="margin-bottom: 5px;">a) A stool specimen positive for a pathogen (such as Salmonella, Shigella, E. coli 0157:H7, Campylobacter species, rotavirus) <li style="margin-bottom: 5px;">b) At least one of the following GI sub-criteria present <ol style="list-style-type: none"> <li style="margin-bottom: 5px;">i. Nausea <li style="margin-bottom: 5px;">ii. Vomiting <li style="margin-bottom: 5px;">iii. Abdominal pain <li style="margin-bottom: 5px;">iv. Diarrhea | <p>Care must be taken to exclude non-infectious causes of symptoms. For instance, new medication may cause diarrhea, nausea/vomiting; initiation of new enteral feeding may be associated with diarrhea; nausea or vomiting may be associated with gallbladder disease.</p> <p>Presence of new GI symptoms in a single resident may prompt enhanced surveillance for additional cases.</p> <p>In the presence of an outbreak, stool from specimens should be sent to confirm the presence of norovirus, or other pathogens (such as rotavirus and E. coli 0157:H7).</p> |

81

Gastrointestinal Tract Infections

Criteria

B. Norovirus gastroenteritis

Both criteria 1 and 2 present

1. At least one of the following GI sub-criteria
 - a) Diarrhea, three or more liquid or watery stools above what is normal for the resident within a 24-hour period
 - b) Vomiting, two or more episodes in a 24-hour period
2. A stool specimen positive for detection of norovirus either by electron microscopy, enzyme immune assay, or by a molecular diagnostic test such as polymerase chain reaction (PCR).

Comments

In the absence of laboratory confirmation, an outbreak (2 or more cases occurring in a LTCF) of acute gastroenteritis due to norovirus infection in a LTCF may be assumed to be present if all of the following criteria are present (“Kaplan criteria”)

- a) Vomiting in more than half of affected persons
- b) A mean (or median) incubation period of 24-48 hours
- c) A mean (or median) duration of illness of 12-60 hours
- d) No bacterial pathogen is identified in stool culture.

82


| Criteria | Comments |
|---|---|
| <p>C. <i>Clostridium difficile</i> gastroenteritis</p> <p>Both criteria 1 and 2 present</p> <p>1. One of the following GI sub-criteria</p> <p>a) Diarrhea, three or more liquid or watery stools above what is normal for the resident within a 24-hour period</p> <p>b) The presence of toxic megacolon (abnormal dilation of the large bowel documented on radiology)</p> <p>2. One of the following diagnostic sub-criteria</p> <p>a) The stool sample yields a positive laboratory test result for <i>C. difficile</i> toxin A or B, or a toxin-producing <i>C. difficile</i> organism is identified in a stool culture or by a molecular diagnostic test such as PCR</p> <p>b) Pseudomembranous colitis is identified during endoscopic examination or surgery, or in histopathologic examination of a biopsy specimen.</p> | <p>A "primary episode" of <i>C. difficile</i> infection (CDI) is defined as one that has occurred without any previous history of CDI, or that has occurred more than 8 weeks after the onset of a previous episode of CDI.</p> <p>A "recurrent episode" of CDI is defined as an episode of CDI that occurs 8 weeks or less after the onset of previous episode, provided the symptoms from the earlier (previous) episode resolved</p> <p>Individuals previously infected with <i>C. difficile</i> may continue to remain colonized even after symptoms resolve</p> <p>In the setting of a GI outbreak, individuals could test positive for <i>C. difficile</i> toxin due to ongoing colonization and be co-infected with another pathogen. It is important that other surveillance criteria are used to differentiate infections in this situation.</p> |

Gastrointestinal Tract Infections

83

CDI LabID Event (different than an infection)

- *C. difficile* positive laboratory assay, tested on a loose-unformed stool specimen, and collected while a resident is receiving care from the LTCF, and the resident has no prior *C. difficile* positive laboratory assay collected in the previous two weeks (<14 days) while receiving care from the LTCF



84

Why is *C. difficile* Surveillance Important?

C. difficile is the leading cause of acute diarrhea in nursing home residents

C. difficile infections contribute to approximately 14,000 deaths/year

~ 90% elderly

Prevention activities, like antimicrobial stewardship programs and hand hygiene are shown to prevent the spread of *C. difficile* and other infections

85

SETTING OF NOROVIRUS OUTBREAKS REPORTED THROUGH THE NATIONAL OUTBREAK REPORTING SYSTEM (NORS), 2009-2012


| Exposure setting | Number of Outbreaks | Percentage of Outbreaks |
|--------------------------------|---------------------|-------------------------|
| Health care facility | 2189 | 62.7% |
| Restaurant or banquet facility | 771 | 22.1% |
| School or day-care facility | 214 | 6.1% |
| Private residence | 69 | 1.9% |
| Other/multiple settings | 251 | 7.2% |

Data on specific settings are restricted to outbreaks with a single exposure setting; for foodborne outbreaks, setting refers to the setting where implicated food was consumed.


<https://www.cdc.gov/norovirus/images/settings-lg.jpg>

Knowledge Check

86



87



88

What type of infection does Ms. Hammer have?

- A. *C. difficile*
- B. Gastroenteritis
- C. Norovirus
- D. Just an upset stomach



89

Gastrointestinal Tract Infections

| Criteria | Comments |
|--|--|
| <p>B. <i>Norovirus gastroenteritis</i></p> <p>Both criteria 1 and 2 present</p> <p>1. At least one of the following GI sub-criteria</p> <ul style="list-style-type: none"> a) Diarrhea, three or more liquid or watery stools above what is normal for the resident within a 24 hour period b) Vomiting, two or more episodes in a 24 hour period <p>2. A stool specimen positive for detection of norovirus either by electron microscopy, enzyme immune assay, or by a molecular diagnostic test such as polymerase chain reaction (PCR).</p> | <p>In the absence of laboratory confirmation, an outbreak (2 or more cases occurring in a LTCF) of acute gastroenteritis due to norovirus infection in a LTCF may be assumed to be present if all of the following criteria are present ("Kaplan criteria")</p> <ul style="list-style-type: none"> a) Vomiting in more than half of affected persons b) A mean (or median) incubation period of 24-48 hours c) A mean (or median) duration of illness of 12-60 hours d) No bacterial pathogen is identified in stool culture. |



90

Gastrointestinal Tract Infections

| Criteria | Comments |
|--|--|
| <p>A. <u>Gastroenteritis</u></p> <p>At least one of the following criteria present</p> <ol style="list-style-type: none"> 1. Diarrhea, three or more liquid or watery stools above what is normal for the resident within a 24-hour period 2. Vomiting, two or more episodes in a 24-hour period 3. Both of the following sign/symptom sub-criteria present: <ol style="list-style-type: none"> a) A stool specimen positive for a pathogen (such as <i>Salmonella</i>, <i>Shigella</i>, <i>E. coli</i> 0157:H7, <i>Campylobacter</i> species, rotavirus) b) At least one of the following GI sub-criteria present <ol style="list-style-type: none"> i. Nausea ii. Vomiting iii. Abdominal pain iv. Diarrhea | <p>Care must be taken to exclude non-infectious causes of symptoms. For instance, new medication may cause diarrhea, nausea/vomiting; initiation of new enteral feeding may be associated with diarrhea; nausea or vomiting may be associated with gallbladder disease.</p> <p>Presence of new GI symptoms in a single resident may prompt enhanced surveillance for additional cases.</p> <p>In the presence of an outbreak, stool from specimens should be sent to confirm the presence of norovirus, or other pathogens (such as rotavirus and <i>E. coli</i> 0157:H7).</p> |

91

Questions?



SPICE 

92