This policy has been adopted by UNC Hospitals for its use in infection control. It is provided to you as information only.

Isolation Precautions

Attachment 7: Infection Control Recommendations for Multiple Patients/Healthcare Personnel with Signs/Symptoms of Gastroenteritis

The following measures will be implemented to prevent the spread of gastroenteritis:

- I. Patients
 - A. Patients with symptoms of gastroenteritis (for example, 3 or more loose stools in 24 hours; vomiting and diarrhea) will be placed on Enteric Precautions. Notify Infection Prevention (984-974-7500 or page using myUNC Health Directory) if additional cases occur.
 - B. Ideally, do not transfer symptomatic patients to an unaffected nursing unit until at least 48 hours after symptoms have ended.
 - C. Consider cohorting-affected patients to designated staff.
 - D. In certain outbreak situations additional enhanced precautions may be necessary (e.g. closure of a unit) at the discretion of Infection Prevention.
- II. Staff
 - A. Staff with symptoms of gastroenteritis will not be allowed to work until asymptomatic for 48 hours.
 - B. Staff will not eat or drink on the unit (consuming meals in the breakroom is allowed).
 - C. Staff will perform hand hygiene with soap and water and in accordance with the <u>Infection Prevention policy: Hand Hygiene and Use of Antiseptics for Skin Preparation</u> Gastroenteritis is spread by the fecal-oral route therefore, hand hygiene is essential before eating and after using the bathroom.
- III. Housekeeping
 - A. Gloves and gown should be worn to clean Enteric Precautions rooms.
 - B. Norovirus
 - 1. Environmental Services will perform a thorough cleaning of the patient's room and floor with a sporicidal cleaning agent (e.g., a 1:10 solution of bleach and water).
 - For outbreaks (e.g., ≥3 patients on a nursing unit overlapping in time), all routine daily cleaning by Environmental Services should be done with a sporicidal cleaning agent for the entire unit, until 48 hours after last patient is symptomatic and there are no further cases among staff or patients.
 - C. C. difficile and Rotavirus
 - 1. Terminal cleaning of the room surfaces and floors will be done using a sporicidal cleaning agent
 - 2. Additional cleaning may be indicated for outbreaks and as directed by Infection Prevention.

For questions about this policy or for other infection control concerns, contact Infection Prevention at 984-974-7500