



DEVELOPMENT OF AN INFECTION CONTROL PROGRAM FOR LONG-TERM CARE FACILITIES

Marty Cooney, DrPH, MPH, ME, BSN, RN, CIC

Associate Director,

NC Statewide Program for Infection Control and Epidemiology
(SPICE)

<https://spice.unc.edu/>

Discuss

Describe

List

Discuss

***Discuss
nursing
home
complex
demographics***

***Describe
unique
infection
prevention
challenges
associated
with LTCFs***

***List factors
contributing
to infections
in the
elderly***


***Discuss
components
of a LTCF
infection
control
program***

Long-term Care Environment

- Long-term care (LTC) generally refers to the large range of facilities that provide care to individual(s) unable to achieve independent self or assisted care:
 - Nursing home
 - Skilled nursing and
 - Assisted living facilities
- Encompasses medical, physical, and psychosocial care
- Typically serve as the resident's home



Figure 20. Percent distribution of post-acute and long-term care services users, by sector and age group: United States, 2017 and 2018



https://www.cdc.gov/nchs/data/series/sr_03/sr03-047.pdf

NATIONAL CENTER FOR HEALTH STATISTICS

Nursing Home Demographics (published May 2022)

- Number of nursing homes: 15,600
- Proportion of nursing homes with for-profit ownership: 70.0%
- Number of licensed beds: 1.7 million
- Number of residents: 1.3 million

https://www.cdc.gov/nchs/data/series/sr_03/sr03-047.pdf

Nursing Homes



- Changed significantly over the past several decades
- Government regulation and consumer pressure
 - Highly regulated
 - Increased acuity of residents
 - Medical needs more complex

“The problem is that nursing homes *still operate on antiquated assumptions made decades ago about the complexity of care their residents require*. Previously, older adults populated nursing homes primarily for custodial care and needed little in the way of medical intervention.

Scientific advances have introduced treatments for illnesses that previously were synonymous with death but now can be managed with medicine and therapies. As a result, those who wind up in nursing homes—many after typically brief hospital stays—are extraordinarily frail, with multiple underlying conditions that demand elaborate medication regimens. “

“there is a notable rise in young patients bringing unique challenges. They are disabled by neurological disorders, trauma, or drug abuse, some have myriad afflictions from birth. younger adults are estimated to be the fastest-growing subpopulation in post-acute and long-term care, increasing 16.5 percent.”

YOUNGER ADULT IN LTC

- LTC is no longer synonymous with “geriatric care”
- Adults aged 31-64 years fastest growing population
- Require different approaches in care



[The Younger Adult in the Long-Term Care Setting | AMDA | The Society for Post-Acute and Long-Term Care Medicine](#)

Discuss

Describe

List

Discuss

*Discuss
nursing home
complex
demographics*

***Describe
unique
infection
prevention
challenges
associated
with LTCFs***

***List factors
contributing
to infections
in the
elderly***

***Discuss
components
of a LTCF
infection
control
program***

INFECTION PREVENTION PROGRAMS

- Infection Prevention and Control (IPC) programs are inadequately staffed, as much as four-fold less than their acute care hospital counterparts
- IPs wear multiple hats
- Less than 10% have specialized training
- Difference in social environment
- Populations differ across LTCFs

*“IP spend less than a third of their time on IP work; only 40% have specialized training; and less than 10% are certified”
...3/29/23*

Council of State and Territorial Epidemiologists (CSTE): Recommendations for Surveillance and Reporting of Healthcare-Associated Infections in Long Term Care Facilities

INFECTION PREVENTIONIST

- Virtual audience:
 - Please put into the chat box how many other roles you are responsible for
 - Please put in the chat box how long you have been in the role of IP
- In-person-
 - Show of hands

STAFFING

- CMS Final Rule effective June 21, 2024: ***“Medicare and Medicaid Programs; Minimum Staffing Standards for Long-Term Care Facilities and Medicaid Institutional Payment Transparency Reporting”***.

- The regulations at § 483.71 must be implemented by August 8, 2024, for all facilities.
- The regulations at § 483.35(b)(1) and (c)(1) must be implemented by May 11, 2026, for non-rural facilities and May 10, 2027, for rural facilities as defined by the Office of Management and Budget.
- The regulations at § 483.35(b)(1)(i) and (ii) must be implemented by May 10, 2027, for non-rural facilities and May 10, 2029, for rural facilities as defined by the Office of Management and Budget.
- The regulations at §§ 438.72(a) and 442.43 must be implemented by all States and territories with Medicaid-certified nursing facilities and intermediate care facilities for individuals with intellectual disabilities beginning May 10, 2028.

cms.gov/newsroom/fact-sheets/medicare-and-medicaid-programs-minimum-staffing-standards-long-term-care-facilities-and-medicaid-0

federalregister.gov/documents/2024/05/10/2024-08273/medicare-and-medicaid-programs-minimum-staffing-standards-for-long-term-care-facilities-and-medicaid

AHCA Files Lawsuit Against Federal Staffing Mandate

Nursing home associations and providers ask court to vacate minimum staffing rule that would reduce access to long term care

WASHINGTON, D.C. – The American Health Care Association (AHCA), joined by the Texas Health Care Association (THCA) and several Texas long term care facilities [filed a lawsuit](#) late Thursday in the Northern District of Texas against the U.S. Department of Health and Human Services (HHS) and the Centers for Medicare and Medicaid Services (CMS) for exceeding their statutory authority and arbitrarily and capriciously issuing the Minimum Staffing Standards for Long-Term Care Facilities final rule. The lawsuit asks the court to issue an order and judgment setting aside the new staffing requirements that were finalized by CMS on May 10.


June 18, 2024, Washington, DC — Following [its early June decision](#) to file to join the American Health Care Association's (AHCA) lawsuit against the Department of Health and Human Services (HHS) and the Centers for Medicare and Medicaid Services (CMS), LeadingAge, the association of nonprofit providers of aging services, including nursing homes, today announces its official co-plaintiff status.

Core Staffing Proposals


1. Minimum nurse staffing standards of 0.55 hours per resident day (HPRD) for RNs and 2.45 HPRD for Nurses Aides (NAs);
2. Requirement to have an RN onsite 24 hours a day, seven days a week; and
3. Enhanced Facility Assessment

Includes staggered implementation schedule and possible hardship exemption

Timeline for Non-rural Facilities



Timeline for Rural Facilities



Permitting Regulatory Flexibility

- LTC facilities may qualify for a temporary hardship exemption from the minimum nurse staffing HPRD standards only if they are able to meet specific criteria demonstrating the following:
 - Workforce unavailability based on location
 - Demonstrate good faith efforts to hire and retain staff
 - A financial commitment to staffing



INFECTION PREVENTION CHALLENGES

- Never been required to deal with emerging infectious diseases
- Regulatory oversight - Isolation should be the least restrictive possible (CMS)
 - PPE used much less frequently
 - Education/monitoring absent or inadequate





JAMDA

journal homepage: www.jamda.com

Original Study

Nursing Home Profit Margins and Citations for Infection Prevention and Control



Hari Sharma PhD*, Lili Xu MS

Department of Health Management and Policy, The University of Iowa, Iowa City, IA, USA

- Results indicated a need for more **clinical guidelines and public policies** designed to bolster providers' limited resources
- A major part of the struggle is the **ongoing staffing shortage** — since IPC measures can be labor-intensive and time-consuming.
- **Supply chain issues** "Policies should ensure ongoing preparedness and oversight that supports NHs organizational readiness to deal with future pandemics. This includes **support** for facility renovations, education and IPC resources such as PPE supplies, testing kits, vaccinations, and other supplies."

Conclusions: Findings indicate that nursing homes may need more resources to prevent citations for infection prevention and control.

<https://www.jamda.com/action/showPdf?pii=S1525-8610%2821%2900333-9>



Discuss

Describe

List

Discuss


*Discuss
nursing home
complex
demographics*

*Describe
unique
infection
prevention
challenges
associated
with LTCFs*

***List factors
contributing
to infections
in the elderly***

***Discuss
components
of a LTCF
infection
control
program***

ELEMENTS REQUIRED FOR AN INFECTION TO OCCUR



Breaking the Chain of Infection



North Carolina
SPICE
Statewide Program for
Infection Control & Epidemiology

Do you know how?

SPICE 



The Links in the Chain of Infection

1. Infectious Agent – The pathogen (e.g., bacteria, virus, fungus, parasite).

✓ **Break it by:** Sterilization, disinfection, and antimicrobial stewardship.

2. Reservoir – Where the pathogen lives and multiplies (e.g., humans, animals, water, soil, medical equipment).

✓ **Break it by:** Cleaning and disinfecting environments, proper waste disposal, and managing contaminated equipment.

3. Portal of Exit – How the pathogen leaves the reservoir (e.g., coughing, sneezing, blood, bodily fluids).

✓ **Break it by:** Covering coughs/sneezes, using PPE, proper wound care, and isolation precautions.



The Links in the Chain of Infection

4. Mode of Transmission – How the pathogen travels (e.g., direct contact, droplets, airborne, fomites).

- **Break it by:** Hand hygiene, social distancing, cleaning surfaces, and using barriers like gloves and masks.

5. Portal of Entry – How the pathogen enters a new host (e.g., mucous membranes, wounds, respiratory tract).

- **Break it by:** Using PPE, maintaining skin integrity, and safe catheter/tube management.

6. Susceptible Host – A person who can get infected (e.g., immunocompromised, elderly, infants).

- **Break it by:** Vaccination, good nutrition, managing chronic conditions, and reducing exposure risks.

HEALTHCARE- ASSOCIATED INFECTIONS (HAI)

- 1 – 3 million serious infections annually
- Infections include:
 - UTI, diarrheal disease, antibiotic-resistant staph infections, and others
- Major cause of hospitalization
- 380,000 die of infections in LTCFs annually



SPICE 

HEALTHCARE- ASSOCIATED INFECTIONS (HAI)

- Account for 26% of all serious adverse events
- **59% deemed preventable**
- Among the most frequent causes of transfer to acute care hospitals and 30-day hospital readmissions.
- Cost of infection-related hospitalizations was estimated to be \$83 million in single month

OIG. Adverse Events in Skilled Nursing Facilities: National Incidence Among Medicare Beneficiaries, OEI-06-11-00370




SPECIFIC INFECTIONS IN LTCFS

- Urinary Tract Infections
 - 30% of hospital readmissions in 30 days
- Respiratory Track Infections
 - Pneumonia 5th lead cause of death >65
- Skin and Soft Tissue
- Gastroenteritis
 - *C difficile* (most common)
- Conjunctivitis
- Bacteremia(s)
 - 50% related to UTI



SPECIFIC INFECTIONS IN LTCFS



[CDC Home](#)[Search](#)[Health Topics A-Z](#)**MMWR***Weekly*

March 11, 2005 / 54(09);220-223

Transmission of Hepatitis B Virus Among Persons Undergoing Blood Glucose Monitoring in Long-Term--Care Facilities --- Mississippi, North Carolina, and Los Angeles County, California, 2003--2004

Regular monitoring of blood glucose levels is an important component of routine diabetes care (*1*). Capillary blood is typically sampled with the use of a fingerstick device and tested with a portable glucometer. Because of outbreaks of hepatitis B virus (HBV) infections associated with glucose monitoring, CDC and the Food and Drug Administration (FDA) have recommended since 1990 that fingerstick devices be restricted to individual use (*2,3*). This report describes three recent outbreaks of HBV infection among residents in long-term--care (LTC) facilities that were attributed to shared devices and other breaks in infection-control practices related to blood glucose monitoring. Findings from these investigations and previous reports suggest that recommendations concerning standard precautions and the reuse of fingerstick devices have not been adhered to or enforced consistently in LTC settings (*2--5*). The findings underscore the need for education, training, adherence to standard precautions, and specific infection-control recommendations targeting diabetes-care procedures in LTC settings (*4--6*) ([Box 1](#)).



RESIDENT FACTORS (*NON-MODIFIABLE*) CONTRIBUTING TO INFECTIONS

- Medications affecting resistance to infection
- Limited physiologic reserve
- Compromised host defenses (↓ cough reflex, thinning skin, decreased tear production and immune dysfunction)
- Coexisting chronic diseases
- Impaired responses to infection
- Increase frequency of therapeutic toxicity




Health needs increase with age among nursing home residents

According to the CDC, top diagnoses among nursing home residents are:

Medical Diagnosis	Percentage of Nursing Home Residents
Hypertension	71.5%
Alzheimer's disease or other dementias	47.8%
Depression	46.3%
Heart Disease	38.1%
Diabetes	32.0%

MODIFIABLE FACTORS CONTRIBUTING TO INFECTION TRANSMISSION

- Lack of a staff member dedicated to the function of infection prevention and control
 - Staff education, **monitoring** and competency
- Semi-private rooms
- Inadequate ventilation systems and/or systems maintenance
- Residents sharing space, air, food in a crowded institutional setting
- Multiple visitors



Discuss

Describe

List

Discuss

*Discuss
nursing home
complex
demographics*

*Describe
unique
infection
prevention
challenges
associated
with LTCFs*

*List factors
contributing
to infections
in the
elderly*

***Discuss
components
of a LTCF
infection
control
program***

REGULATORY AND/OR ACCREDITING AGENCIES

- OSHA (Occupational Safety and Health Administration)
- CMS
- TJC (The Joint Commission)



ADDITIONAL NC STATE REGULATIONS

- Rules Governing the Sanitation of Hospitals, Nursing and Rest Homes, Sanitariums, Sanitoriums and Other Institutions - 15A NCAC 18A .1300
- NC Communicable Disease Rule 10A NCAC 41A .0206.
- NC Rules for the Licensing of Nursing Homes and Beds in Homes for the Aged Licensed as Part of a Nursing Home

SHEA/APIC GUIDELINES:

Infection prevention and control in the long-term care facility

- *In this document, as in several published HICPAC, SHEA, and APIC guidelines, each recommendation is categorized based on existing scientific evidence, theoretical rationale, applicability, and national or state regulations*

*Healthcare Infection Control Practices Advisory Committee (HICPAC)

*Society Healthcare Epidemiology of America (SHEA)

*Association for Professionals in Infection Control and Epidemiology (APIC)

Smith et al; AJIC September 2008

CATEGORIZATION OF RECOMMENDATIONS

- Category IA: Strongly recommended and strongly supported
- Category IB: Strongly recommended with some support
- **Category IC: Required by law/regulation**
- Category II: Recommended for implementation
- No Recommendation: Unresolved issues

LTCF INFECTION PREVENTION PROGRAM

- An active, effective, facility-wide infection prevention program should be established in the LTCF (Cat 1C).
 - The Purpose of the program is to reduce the risk of development and spread of infectious disease
- The IP Program must comply with federal, state and local regulations (Cat 1C)


INFECTION PREVENTION AND CONTROL PROGRAM (IPCP)

- **§483.80 Infection Control**
 - The facility must establish and maintain an infection **prevention and control** program designed to provide a safe, sanitary and comfortable environment and to help prevent the development and transmission of **communicable** disease and infection

INFECTION PREVENTION AND CONTROL PROGRAM (IPCP)

- Requires system for preventing, identifying, reporting, investigating and controlling infections and communicable diseases that:
 - Covers all residents, staff (*direct and indirect care*), visitors, volunteers and other service providers. Expectation that facilities tailor the emphasis of their IPCP for visitors and to work to prevent transmission
For example, “*screening may be* passive using signs to alert *family members and* visitors with signs and symptoms of communicable diseases not to enter. *More active screening may include the completion of a screening tool or questionnaire which elicits information related to recent exposures or current symptoms. That information is reviewed by the facility staff and the visitor is either permitted to visit or is excluded*”
 - **Is based on the individual facility assessment**
 - Follows accepted national standards

FACILITY-WIDE ASSESSMENT




- “Determine what resources are necessary to care for its residents competently during both day-to-day operations (*including nights and weekends*) and emergencies”
- “The facility must review and update that assessment:
 - As necessary
 - At least annually
 - Whenever there is, or facility plans for, any change that would require a substantial modification to any part of this assessment”
- Must include a facility-based and community-based risk assessment (*MDROs, HAIs and communicable diseases*)

Source: Appendix PP State Operations manual_8/24

Strengthening the Facility Assessment

- Clarifying that facilities must use **evidence-based methods** when care planning for their residents, including consideration for those residents with behavioral health needs;
- Requiring that facilities use the facility assessment to **assess the specific needs of each resident** in the facility and to adjust as necessary based on any significant changes in the resident population;
- Requiring that facilities **include the input of facility staff**, including, but not limited to, nursing home leadership, management, direct care staff (i.e., nurse staff), representatives of direct care staff, and staff who provide other services; and,
- Requiring facilities to develop a staffing plan to maximize recruitment and retention.



<https://qioprogram.org/tools-resources/facility-assessment-tool>

- The facility's resident population (*the number and facilities capacity*)
- The care required for resident population:
 - Types of diseases, conditions, physical and *behavioral health needs*, cognitive disabilities; Overall acuity; Other pertinent facts that are present within that population
- Staff competencies and *skill set* necessary to provide level of care
- Physical environment, equipment, services
- Ethnic, cultural or religious factors
- *Must use to :*
 - *Inform staffing decisions; Consider specific staffing needs for each resident unit; Consider specific staffing needs for each shift; Develop and maintain a plan to maximize recruitment and retention*

INFECTION PREVENTION RISK ASSESSMENT



spice.unc.edu/resources/spice-ltc-infection-prevention-risk-assessment/
cdc.gov/long-term-care-facilities/hcp/training/index.html



INFECTION CONTROL RISK ASSESSMENT

*Infection Control
Risk Assessment*



Priorities



Goals



Infection Control Plan

EVENT	PROBABILITY OF OCCURRENCE <i>(How likely is this to occur)¹</i>				RISK LEVEL OF FAILURE <i>(What would be the most likely)²</i>				POTENTIAL CHANGE IN CARE <i>(Will treatment/care be needed for resident/staff)³</i>				PREPAREDNESS <i>(Are processes in place and can they work)⁴</i>			YEAR: _____ RISK LEVEL Add rankings (score of 8 or > are considered highest priority for improvement efforts)
	High	Med	Low	None	Life Threatening	Permanent Harm	Temp Harm	None	High	Med	Low	None	Poor	Fair	Good	
Score	3	2	1	0	3	2	1	0	3	2	1	0	3	2	1	
Example: Lack of Communication with Transferring Facility	2				1				2				1			6
External Factors (Community, Demographics)																
Identify other risk factors in the community based on geographic location (coast, mountains etc.)																
Risk of TB in the community																
Risk of emerging infectious disease in the community																
Internal Factors (Facility Related)																
Facility Associated Infection(s)																
Symptomatic urinary tract infection (SUTI)																
Influenza like illness																
Pneumonia																

spice.unc.edu/resources/spice-ltc-infection-prevention-risk-assessment/



2 TYPES OF EVENTS/RISKS



- **Community/External**

- TB risk (HCP & residents)
- **COVID transmission in community**
- Geographical area & environmental issues such as flooding, mudslides, hurricane, tornado, legionella, etc.
- Population served & socioeconomic status such as retirement community, rural, low income, drug abuse, etc.
- Regulatory – CMS, DHSR, OSHA

- **Facility specific/Internal**

- Healthcare associated infections
- Antibiotic stewardship/ MDROs
- Exposure related events
- HCP compliance
- Resident/family
- New services/construction
- Procedures/devices

FOUR FACTORS TO CONSIDER: RANKING THE RISK

Probability of Occurrence (Likelihood)

- High: If there were more events than baseline numbers or more than experienced historically
- Medium: If there were a similar number of events experienced historically
- Low: If there were fewer events than expected or experienced historically
- None: No events occurred

Risk Level of Failure

- Life-Threatening: Event associated with high rates of mortality
- Permanent Harm: Event associated with loss of limb or permanent change in status
- Temporary Harm: Event associated with a temporary change in ambulation

Potential Change in Care

- High: Event resulted in transfer to higher level of care (hospital)
- Medium: Event resulted in major change to resident's care plan (acquisition of *C difficile* for example)
- Low: Event resulted in minor/short term modification to treatment (change in VS routine for example)
- None: No change in treatment or care plan

Preparedness

- Poor: No policies or procedures or process in place
- Fair: Policies/procedures in place but no monitoring to ensure compliance
- Good: Policies/procedures in place and compliance being monitored with staff feedback

INFECTION PREVENTION AND CONTROL PROGRAM (IPCP)

- Must include, at a minimum policies and procedures that address - **§483.80(a)(2)**:
 - Surveillance (communicable diseases **and** infections)
 - Reporting
 - Standard and Transmission-based Precautions (**define and explain application and how to utilize**)
 - Emphasis that isolation should be the least restrictive
- Ensure staff are aware of policies
- Annual review of the IPCP and update as needed

Appendix PP State Operations manual 8/24

POLICIES INCLUDE

- Hand hygiene
 - *ABHR preferential use*
- Selection and use of PPE
- Addressing use of facemasks for residents with new respiratory symptoms
- Addressing resident room assignment
- How to manage when on TBP and single room not available
- Limiting movement if on TBP

POLICIES INCLUDE-*continued*

- Respiratory Hygiene/Cough Etiquette
 - Increase prevalence of respiratory infections **should** have facemasks available and **offer** them to visitors and **others entering the facility**.
 - Post signs with instructions **on visitation restriction for those with symptoms**
- Environmental cleaning and disinfection
 - Routine cleaning and disinfection/frequently touched surfaces
 - Privacy curtains-changed when visibly dirty
 - Shared equipment-routine cleaning and disinfection
 - Objective methods for evaluation
 - Direct observation; Fluorescent markers; Adenosine triphosphate (ATP)

Appendix PP State Operations manual 10/22- 2/23, 8/24

POLICIES INCLUDE-*continued*

- Occupational Health
 - Work restrictions, prohibiting contact with food or residents
 - Assess risk for TB based on **exposure or cases of TB in the facility** and screen
 - Monitor for clusters or outbreaks among staff
 - Exposure control plan

Educate staff, residents and visitors on the IPCP
Monitor adherence

POLICIES INCLUDE-*continued*

- Linens
 - Use standard precautions if potentially contaminated (e.g., gloves, **gowns when sorting and rinsing**)
 - No special precautions (e.g., double bagging, **melting bags**) or categorizing (**e.g., biohazard, color-coded**) for linen originating in transmission-based precaution rooms is necessary



Appendix PP State Operations manual 10/22; 2/23; 8/24

Legionellosis

- “*Legionellosis*” refers to two clinically and epidemiologically distinct illnesses:
 - **Legionnaires’** disease, which is typically characterized by fever, myalgia, cough, and clinical or radiographic pneumonia and
 - **Pontiac fever**, a milder illness without pneumonia (e.g., fever and muscle aches).

Caused by Legionella bacteria


Water Management

- *Legionella can grow in parts of building water systems that are continually wet (e.g., pipes, faucets, water storage tanks, decorative fountains), and certain devices can spread contaminated water droplets via aerosolization (i.e., converted into a spray or mist in the air).*
- *Facilities must be able to demonstrate its measure to minimize the risk of Legionella and other opportunistic pathogens in building water systems such as by having a documented water management plan.*
 - *An assessment (identify where could grow and spread*
 - *Measures to prevent growth and how to monitor them*

CMS does not require water cultures as part of routine program validation

INFECTION PREVENTION AND CONTROL PROGRAM (IPCP) **F881**

- An antibiotic stewardship program that includes antibiotic use protocols and a system to monitor use
- A system for recording incidents identified and the corrective actions taken



Antibiotic Stewardship Program

- Incorporate monitoring of antibiotic use, including the frequency of monitoring/review. Monitor/review *response to antibiotics, and laboratory results when available, to determine if the antibiotic is still indicated or adjustments should be made (e.g., antibiotic time-out)*
- *Facilities should provide* feedback (e.g., verbal, written note in record) to prescribing practitioners regarding antibiotic resistance data, their antibiotic use and their compliance with facility antibiotic use protocols *to improve prescribing practices and resident outcomes.*
- *Require antibiotic orders to include the indication, dose, and duration.*

ADMINISTRATIVE STRUCTURE


(Committee)

- Oversight of the IP program should be defined and should include participation of the IP, administration, nursing staff, and physician staff (Category II)
 - Meet on regular basis
 - Written minutes with action plans and recommendations
 - Evaluate effectiveness
 - Review of IP data
 - Approve policies and procedures



INFECTION PREVENTIONIST-F882

- Collection and analysis of infection data
- Evaluation of products and procedures
- Development of policies
- Consultation
- Education



- Implementation of mandated changes
- Application of epidemiologic principles-*outbreak management*
- Antimicrobial management
- Research
- High quality services in a cost-efficient manner

483.80(b) Infection Preventionist

*Issued 10-21-22; Effective 10-21-22; Implementation 10-24-22; Revised 2/23;
8/24*

- The facility must designate one or more individuals as the infection preventionist (IP) who is responsible for assessing, developing, implementing, monitoring, and managing the IPCP.
- The IPCP includes content required in §§483.80(a)(1)-(4), (F880, Infection Prevention and Control and at F881, Antibiotic Stewardship Program (ASP)). While an ASP is a team effort, the IP is responsible for ensuring the program meets the requirements for ASPs (at §483.80(a)(3), F881).
- The IP should review and approve infection prevention and control training topics and content, as well as ensure facility staff are trained on the IPCP (for further information, see §483.95(e), F945, Infection Control Training).
 - Does not have to perform the IPCP training, since some facilities may have designated staff development personnel

- **Have primary professional training:**

- Nursing-earned a certificate/diploma or degree in nursing,
- Medical technology-an associate's degree in medical technology or clinical laboratory science,
- Microbiology-earned a bachelor's degree in microbiology,
- Epidemiology-earned a bachelor's degree in epidemiology
- Examples of other related fields of training that are appropriate include physicians, pharmacists and physician's assistants



- **Qualified by education, training, experience or certification**

- *The IP should remain current with infection prevention and control issues and be aware of national organizations' guidelines as well as those from national/state/local public health authorities (e.g., emerging pathogens). **The facility should ensure the individual selected as the IP has the background and ability to fully carry out the requirements of the IP based on the needs of the resident population, such as interpreting clinical and laboratory data.***



Reading Materials

Online embedded reader plus optional printed books

- **MODULE 1:** Long-Term Care Settings
- **MODULE 2:** Microbiology, Epidemiology, and Normal Aging Processes
- **MODULE 3:** The Infection Prevention and Control Program
- **MODULE 4:** Surveillance of Communicable Diseases
- **MODULE 5:** Prevention and Control of Communicable Diseases
- **MODULE 6:** Infection Prevention for Ancillary Services



Online Study Tools:

- **Assessment**-helps determine your current knowledge/areas for improvement
- **Module Quizzes**-need further study
- **Flashcards**-Practice key terms and concepts
- **Glossary**-Look up key terms
- **Resource Center**-Links to IP information referenced in printed materials, downloadable e-book versions
- **Post-Test**
- **Reports**-Track your progress, compare results of quizzes and pre- and post-tests

<https://learnipc.apic.org/ltc-cip-certification/>



- **Works at least part time in the facility**

- Designated IP hours per week can vary based on the facility and its resident population. Therefore, the amount of time required to fulfill the role must be at least part-time and should be determined by the facility assessment, conducted according to §483.70(e), to determine the resources it needs for its IPCP, and ensure that those resources are provided for the IPCP to be effective.
- Based upon the assessment, facilities should determine if the individual functioning as the IP should be dedicated solely to the IPCP. *A facility should consider resident census as well as resident characteristics, types of units such as respiratory care units, memory care, skilled nursing and the complexity of the healthcare services it offers as well as outbreaks and seasonality of infections such as influenza in determining the amount of IP hours needed.*
- The IP must have the time necessary to properly assess, develop, implement, monitor, and manage the IPCP for the facility, address training requirements, and participate in required committees such as QAA.
- *Must physically work onsite in the facility*

- **Completed specialized training**

- *An IP must have obtained specialized IPC training beyond initial professional training or education prior to assuming the role. Training can occur through more than one course, but the IP must provide evidence of training through a certificate(s) of completion or equivalent documentation.*

- *Infection prevention and control program overview*
 - *The infection preventionist's role*
 - *Infection surveillance*
 - *Outbreaks*
 - *Principles of standard precautions*
 - *Principles of transmission-based precautions*
 - *Resident care activities (e.g., use and care of indwelling urinary and central venous catheters, wound management, and point-of-care blood testing)*
 - *Preventing respiratory infections (e.g., influenza, pneumonia)*
 - *Tuberculosis prevention*
 - *Occupational health considerations (e.g., employee vaccinations, exposure control plan, and work exclusions)*
 - *Quality assurance and performance improvement*
 - *Antibiotic stewardship*
 - *Care Transitions*
 - *Water management*
 - *Linen management*

Office of Inspection General (OIG): Certain for-profit NH may not have complied with federal requirements regarding the infection prevention position(8/24)

- 76% (76/100) of the for-profit nursing homes complied with Federal requirements pertaining to IPs.
- 17% potentially did not comply with the requirement related to IP completing specialized IPC training **prior** to assuming the role and
- 7 potentially did not comply with the requirement to designate an IP

Figure 1: The Locations of the 100 Sampled Nursing Homes



SPICE Mentorship Program




To request additional information or for questions please contact Marty Cooney at marty_cooney@med.unc.edu

- In collaboration with the North Carolina Department of Health and Human Services (NC DHHS), the North Carolina Statewide Program for Infection Control and Epidemiology (SPICE) would like to encourage your participation in a ***free performance improvement project*** focusing on Infection Prevention and Control (IPC).
 - Funded by CDC via contract with NCDHHS
 - Onsite mentoring
 - Visits, standardized plan of activities and topics



INFECTION PREVENTIONIST (IP)

- The individual designated as the IP (or at least one if there is more than one) must be:
 - A member of the facilities quality assessment and assurance committee
 - Report on the IPCP on a regular basis



<https://www.cms.gov/Medicare/Provider-Enrollment-and-certification/QAPI/qapiresources>



INFECTION PREVENTIONIST (IP)



at a Glance:

A Step by Step Guide to Implementing Quality Assurance and Performance Improvement (QAPI) in Your Nursing Home

- *Reporting may include, but is not limited to, facility process and outcome surveillance, outbreaks (ongoing and any since the last meeting) and control measures, occupational health communicable disease illnesses (e.g., TB, influenza) and the Antibiotic Stewardship Program (ASP) related to antibiotic use and resistance data.*
- *In order to be considered an active participant, the IP should attend each QAA meeting. If the IP cannot attend, another staff member should report on the IP's behalf, but this does not change or absolve the IP's responsibility to fulfill the role of QAA committee member or reporting on the IPCP.*

QUALITY ASSURANCE PERFORMANCE IMPROVEMENT

- Develop, implement and maintain an effective, comprehensive, data-driven QAPI program
 - Address all systems of care and management practices
 - Include clinical care, quality of life and resident choice
 - Define and measure indicators of quality and facility goals
 - Reflect the complexities, unique care and services the facility provides

“Quality Assurance and Performance Improvement (QAPI)” is the coordinated application of two mutually-reinforcing aspects of a quality management system: Quality Assurance (QA) and Performance Improvement (PI). QAPI takes a systematic, interdisciplinary, comprehensive, and data-driven approach to maintaining and improving safety and quality in nursing homes while involving residents and families in practical and creative problem solving.

MONITORING PERFORMANCE: AUDITS

- Quality audits are performed to verify conformance to standards through objective review.
- Should be an opportunity for improvement and not punitive
- Audits can assist the facility in:
 - Establishing a baseline of performance for each activity
 - Identifying what needs to be improved, and
 - Targeting educational needs





"It's not that I don't love you.
It's just that a ten year-old needs his freedom."

- **Timely**
- **Based on data that is valid**
- **Comparisons between peers may be helpful**
- **Sustained**

Feedback is one of many data sources which provide valuable information the facility must incorporate into an effective QAPI program. Each facility must establish and implement written policies and procedures for feedback.

Examples of mechanisms for obtaining resident and staff feedback may include, but are not limited to:

- *Satisfaction surveys and questionnaires;*
- *Routine meetings, e.g., care plan meetings, resident council, safety team, town hall; and*
- *Suggestion or comment boxes*



The screenshot displays the NCDHHS (North Carolina Department of Health and Human Services) Epidemiology website. The main navigation bar includes links for Individuals & Families, Local Health Depts, Healthcare Providers, Schools, Businesses & Community Groups, and Facts & Figures. The left sidebar contains a 'Communicable Disease HOME' section with links to A-Z Diseases & Topics, Tuberculosis, Info for Healthcare Providers, TB Control in N.C., TB Facts & Figures, Programs & Services, Surveillance & Reporting, Disease Laws & Rules, About Us, Contact Us, and Quick Links. The main content area is titled 'Tuberculosis Info for Healthcare Providers' and includes sections for Alerts & Hot Topics, Manuals and Guides, and DHHS patient education materials. The Alerts & Hot Topics section lists four DHHS memos regarding Isoniazid shortages, TB drug shortages, and Tuberculin shortages. The Manuals and Guides section lists the N.C. Tuberculosis Policy Manual, N.C. Guidelines for the Use of Interferon Gamma Release Assays (IGRAs) in TB Diagnosis, and N.C. Communicable Disease Manual. The DHHS patient education materials section lists various brochures and manuals, including Home Isolation Instructions, Information about Isoniazid, Rifampin, and Rifapentine, Instructions for Collecting Sputum for TB, TB and You, You Could Have TB, and Order DHHS TB Control Materials.

NC TB RESOURCES & TB INFECTION CONTROL RISK ASSESSMENT

<https://epi.dph.ncdhhs.gov/cd/tb/providers.html>



TB Screening, Testing and Treatment of U.S. Health Care Personnel (CDC Recommendations 2019)

- U.S. healthcare personnel should be screened for TB upon hire (i.e., preplacement)
- TB screening includes a process that includes:
 - A baseline individual TB risk assessment (2019 updated recommendations)
 - TB symptom evaluation
 - A TB test (e.g., TB blood test or a TB skin test) and
 - Additional evaluation for TB diseased as needed



Figure 3.1 Health care worker collecting a blood




NORTH CAROLINA SPECIFIC RULES

S. Quick Reference for Tuberculin Skin Testing Requirements:

1. Tuberculin Skin Testing (TST) or IGRA (Interferon Gamma Release Assays) testing is **required** by communicable disease/TB rules for:

- **household and other close contacts** of active cases of pulmonary and laryngeal tuberculosis
By: 10 A NCAC 41A .0205
Frequency: at the time of exposure and 3 months post exposure
- persons reasonably **suspected of having tuberculosis disease**
By: 10 A NCAC 41A .0205
Frequency: when suspected
- **inmates in the custody of the Department of Corrections**
By: 10 A NCAC 41A .0205; DOC policy
Frequency: upon incarceration and annually
- **Department of Correction employees with direct inmate contact**
By: 10A NCAC 41A .0205; OSHA; DOC policy
Frequency: upon employment
- **patients in long term care facilities**
By: 10A NCAC 41A .0205; 10A NCAC 13D .2202 & .2209
Frequency: upon admission (two-step for TST or IGRA) & by risk assessment (DFS regulations require an annual screening which can be accomplished by a verbal elicitation of symptoms)
- **long term care facility employees**
By: 10A NCAC 41A .0205; 10A NCAC 13D .2202 & .2209; OSHA
Frequency: upon employment (two-step for TST or IGRA) & by risk assessment (DFS regulations require an annual screening which can be accomplished by a verbal elicitation of symptoms)
- **employees of adult day care centers providing care for persons with HIV infection or AIDS**





NORTH CAROLINA SPECIFIC RULES

10A NCAC 41A.0205

- A 2-step TST or IGRA must be performed on all new residents.

Exceptions

- If the resident is being admitted directly from another hospital, licensed nursing home/adult care home in NC **AND** there is documentation of a 2-step skin test or single IGRA test
→ **NO need to re-test**
- A single TST or IGRA in the following situations
 - Person has ever had a 2-step skin test
 - Person has had a single skin test within the last twelve months

§483.80(d) INFLUENZA AND PNEUMOCOCCAL IMMUNIZATIONS

- Influenza: Facility must develop policies and procedures to ensure that:
 - Before offering, education provided
 - Offered when it becomes available each season is most effective versus date specific (October 1-March 31) annually
 - Right to refuse
 - Documentation
- Pneumococcal disease: Facility must develop policies and procedures to ensure that:
 - Before offering, education provided
 - Offered unless already immunized or medically contraindicated (**CDC has lowered age to 50;10/24**)
 - Right to refuse
 - Documentation

Self-reported doses of the vaccines included above are acceptable

CDC Vaccine Recommendations

(Last webpage update on August 7, 2025)

Legend

Recommended vaccination for adults who meet age requirement, lack documentation of vaccination, or lack evidence of immunity

Recommended vaccination for adults with an additional risk factor or another indication

Recommended vaccination based on shared clinical decision-making

No Guidance/Not Applicable

Vaccine	19-26 years	27-49 years	50-64 years	≥65 years
COVID-19 ⓘ	1 or more doses of 2025–2026 vaccine (See Notes)			2 or more doses of 2025–2026 vaccine (See Notes)
Influenza inactivated (IIV3, cdlIV3) Influenza recombinant (RIV3) ⓘ	1 dose annually			1 dose annually (HD–IIV3, RIV3, or aIIV3 preferred)
Influenza inactivated (aIIV3; HD–IIV3) Influenza recombinant (RIV3) ⓘ	Solid organ transplant (See Notes)			
Influenza live, attenuated (LAIV3) ⓘ	1 dose annually			
Respiratory Syncytial Virus (RSV) ⓘ	Seasonal administration during pregnancy. (See Notes)			60 through 74 years (See Notes) ≥75 years
Tetanus, diphtheria, pertussis (Tdap or Td) ⓘ	1 dose Tdap each pregnancy; 1 dose Td/Tdap for wound management (See Notes)			
	1 dose Tdap, then Td or Tdap booster every 10 years			
Measles, mumps, rubella (MMR) ⓘ	1 or 2 doses depending on indication (if born in 1957 or later)			For health care personnel. (See Notes)
Varicella (VAR) ⓘ	2 doses (if born in 1980 or later)			2 doses
Zoster recombinant (RZV) ⓘ	2 doses for immunocompromising conditions (See Notes)			2 doses
Human papillomavirus (HPV) ⓘ	2 or 3 doses depending on age at initial vaccination or condition	27 through 45 years		
Pneumococcal (PCV15, PCV20, PCV21, PPSV23) ⓘ				See Notes See Notes ⬆

cdc.gov/vaccines/hcp/imz-schedules/adult-age.html



IMPORTANT



Flu, RSV and COVID-19 Vaccines



Vaccines are the best way to protect yourself, your family and your community from severe illness. They help prevent hospital stays and can save lives.

Some people are at higher risk, including:

- Children under 5
- Adults 65 and older
- Pregnant people
- People with certain health conditions, like asthma or diabetes

[Version en Español](#)

covid19.ncdhhs.gov/vaccines

COVID-19 Vaccine Access Update in North Carolina

Governor Josh Stein directed NCDHHS to issue a new Standing Order that makes it easier for adults in NC to get their COVID-19 vaccines at pharmacies. Now, adults 65 and older — and adults ages 18 to 64 with certain health conditions that put them at higher risk — can get vaccinated at a pharmacy without a prescription. The order itself stands in place of a prescription from a health care provider.



https://covid19.ncdhhs.gov/vaccines/frequently-asked-questions-about-covid-19-vaccine

Site review in progress
We are currently reviewing the site to ensure it is up to date for the 2025-26 respiratory virus season. Please check back for updates.

An official website of the State of North Carolina [How you know](#)

NCDHHS NCDPH NC.GOV Select Language

NCDHHS
COVID-19 Response

About Dashboards Vaccines Testing Treatments Guidance Info En Español

Home > Vaccines > Frequently Asked Questions about the COVID-19 Vaccine

Frequently Asked Questions about the COVID-19 Vaccine

What to Know About Standing Orders

What is a Standing Order

Why does NC need a Standing Order

Getting the Vaccine

Who can get the COVID-19 vaccine at a pharmacy without a prescription right

[COVID-19 Vaccine Information](#)

[Frequently Asked Questions about the COVID-19 Vaccine](#)

[Healthier Together - Health Equity Action Network](#)

[Info for Health Care Providers](#)

covid19.ncdhhs.gov/vaccines/frequently-asked-questions-about-covid-19-vaccine

 DEPARTMENT OF HEALTH AND HUMAN SERVICES

Centers for Medicare & Medicaid Services

LTC part starts on page
184

42 CFR Parts 424, 483, and 484

[CMS-1803-P]

RIN 0938-AV28

**Medicare Program; Calendar Year (CY) 2025 Home Health Prospective Payment System
(HH PPS) Rate Update; HH Quality Reporting Program Requirements; HH Value-Based
Purchasing Expanded Model Requirements; Home Intravenous Immune Globulin (IVIG)
Items and Services Rate Update; and Other Medicare Policies**

- **§ 483.80 Infection control.**
 - (g) Respiratory illness reporting--(1) Ongoing reporting.
The facility must electronically report information on
acute respiratory illnesses, including influenza, SARS-CoV-
2/COVID-19, and RSV.

KEY ELEMENTS – EMPLOYEE HEALTH

Immunize

Immunize against vaccine-preventable diseases

- Hepatitis B
- Influenza
- MMR
- Varicella
- Tetanus, diphtheria, pertussis
- [COVID-19](#)

Establish

Establish sick leave policies that encourage:

- Healthcare personnel to stay home when they are ill
- Reporting of signs, symptoms, and diagnosed illnesses that may represent a risk to their patients and coworkers

Adhere

Adhere to federal and state standards and directives applicable to protecting healthcare workers against transmission of infectious agents

EDUCATION AND TRAINING OF HEALTHCARE PERSONNEL ON INFECTION PREVENTION

- Training should be:
 - Job-specific and adapted to the individual healthcare personnel
 - Performed before duties can be assigned and at least annually
 - Additional training to recognized lapses in adherence
 - **Require HCP to demonstrate competency following each training**
 - System of documentation of competency for each healthcare personnel



HEALTHCARE WORKER EDUCATION

➤ Topics should include, but are not limited to:

- Routes of disease transmission
- Hand Hygiene
- Sanitation procedures
- MDROs
- Transmission-based precautions
- OSHA required education



PATIENT, FAMILY AND CAREGIVER INFECTION PREVENTION EDUCATION

- Include information about . . .
 - How infections spread
 - How they can be prevented
 - What signs or symptoms should prompt reevaluation and notification of the patient's healthcare provider
- Instructional materials and delivery should address varied levels of education, language comprehension, and cultural diversity
- Provide education to patients, family members, visitors, and their caregivers



[https://apic.org/Resource_/TinyMceFileManager/IPandYou/IPandYou InfographicPoster 2013.pdf](https://apic.org/Resource_/TinyMceFileManager/IPandYou/IPandYou%20InfographicPoster%202013.pdf)

IN CONCLUSION

- ✓ One person, the IP, should be assigned the responsibility of directing, infection control activities in LTCF.
- ✓ The IP should have a written job description of infection control activities.
- ✓ The IP requires the support of administration in order to function effectively.
- ✓ The IP needs to be guaranteed sufficient time to direct the infection control program.
- ✓ The IP should have written authority to institute infection control measures.

