



**Urinary Tract Infections:  
 Appropriate Prevention, Diagnosis, Treatment and Care**  
 Infection Management AND Antibiotic Stewardship

Mallory McClester Brown, MD  
 University of North Carolina  
 Associate Professor of Family Medicine

1

## Conflict of interest Disclosures

- ▶ The views and opinions expressed in this course are my own and do not reflect the official policy or position of any agency of the U.S. or NC government or UNC.
- ▶ I have NO financial relationships with manufacturer(s) and/or provider(s) of commercial services discussed in this activity.
- ▶ These slides contain materials from a variety of colleagues including CDC, WHO, AHRQ, etc.



2

## Outline of today's session

1. Define urinary tract infection (UTI)
2. Discuss the prevention of UTIs
3. Review the purpose of Urinalysis (UA) and components of UA
4. Review the McGeer Criteria
5. Discuss treatment for UTIs



## A Common Case

- ▶ 84 yo F living in your facility is “sleepier” today per son’s report. Staff note she has eaten less than usual and does seem more fatigued but has no other symptoms or complaints.
- ▶ Son insists that this is what happens “every time Mom has a urine infection” and requests that antibiotic treatment be initiated now. How do you respond?


## CDC NHSN UTI Definitions

- ▶ Urinary Tract Infection (UTI)/Cystitis
  - ▶ infection of the bladder (lower urinary tract).
  
- ▶ Pyelonephritis –
  - ▶ infection of the upper urinary tract (ureters / renal collecting system / kidneys).



5

### Symptoms of Urinary Tract Infection (UTI)



6

## CDC NH UTI Definitions/Points to Clarify



*"Mixed flora" is not considered an organism and cannot be reported.*



*Yeast cannot be reported as an organism for a UTI.*



7

## UTIs: Why do we worry?

Primary cause of bacteremia in LTC residents is due to UTIs!

Incidence of **symptomatic UTIs** in elderly in LTC around 10%

Asymptomatic bacteriuria prevalence:  
30% F/ 10% M



8

## Prevention of UTI



9

## Physiologic Risk Factors for UTIs in the Elderly

- ▶ Physiologic changes of the bladder with aging:
  - ▶ Women
  - ▶ Men



10

## Physiologic Risk Factors for UTIs in the Elderly


### PHYSIOLOGIC CHANGES OF BLADDER WITH AGING:

#### Women:

- ▶ Elevation of vaginal pH due to estrogen deficiency
- ▶ Results in increased ability of bacteria to adhere to the mucosal cells of the bladder.

#### Men:

- ▶ Decreased bactericidal activity of prostatic secretions
- ▶ Increased post-void residual volume of urine
  - Cystocele/rectocele
  - Prostate hypertrophy
  - Neurogenic bladder from comorbidity



11

## Environmental Risk Factors for UTI in the Elderly

### Environmental Risk Factors

- ▶ Indwelling urinary catheters
- ▶ Congregate living
  - Mechanical/chemical restraints
  - Increased exposure to antibiotics
  - Poor infection control techniques



**The more impaired or frail the greater the risk of UTI!**



12

## Physiologic Risk Factors for UTIs in the Elderly (2)

### Functional / Cognitive Impairment

- ▶ Decrease self care
- ▶ Decrease cues to void
- ▶ Increased incontinence and perineal soiling
- ▶ Difficulty finding bathroom / suitable location to void



13

### Risk Factors for CAUTI

Urinary stasis- no below the bladder drainage

Over-distention and pyelonephritis– kinks with backflow

Urethral trauma – catheter tugging

Improper handling of urine collection bag

Duration of catheter use – biofilm buildup

- 5% risk per day of catheterization, >30 days universal asymptomatic bacteriuria



14

**From this information, what are targeted ways to prevent UTI?**



15

## Prevention of UTIs

Hand Hygiene – both residents and staff

Adequate hydration – 30cc/kg of body weight/day


Perineal hygiene after toileting

Routine toileting


*Removing urinary catheter as early as possible.*



16



17



18

## Dipstick Urinalysis

- ▶ Leukocyte esterase
- ▶ Nitrites
- ▶ Protein
- ▶ Blood



19



## Dipstick Urinalysis

- ▶ Leukocyte esterase positive (pyuria)
- ▶ Nitrites: positive (bacteriuria)
- ▶ Protein: small amount may be present
- ▶ Blood: small amount may be present

Leukocyte positive: 50–75% specific; 80-90% sensitive

Pyuria alone not an indication for treatment.



20



## UA: Hematuria

- ▶ Blood is not common with UTIs in older adults.
- ▶ Frank hematuria should be evaluated promptly!
- ▶ Causes:
  - ▶ Stones
  - ▶ Cancer
  - ▶ Trauma
  - ▶ Infection
  - ▶ Hemorrhage.



21



## CDC NHSN UTI Definitions

- ▶ Urinary Tract Infection (UTI)/Cystitis
  - ▶ infection of the bladder (lower urinary tract).
- ▶ Pyelonephritis –
  - ▶ infection of the upper urinary tract (ureters / renal collecting system / kidneys).

22



## Diagnostic Dilemmas for Older Adults with UTI

Common symptoms

Atypical symptoms

Fever?

Getting the history



23

## Evaluation of Possible UTI

- ▶ Vital signs
- ▶ Fever?
- ▶ History and examination
  
- ▶ U/A and C&S *BEFORE* starting antibiotics
- ▶ Clean catch vs I&O catheterization.



24


## Microbiology of UTI

▶ 80% are caused by **gram negative bacilli**

- E.coli, Klebsiella, Enterobacter, Proteus, and Serratia

Few Gram positive bacilli

- Staphylococcus



SPICE 

25

## Urine Culture



This Photo by Unknown Author is licensed under CC BY-NC-ND

- ▶ **Gold STANDARD** to guide appropriate treatment
- ▶ Results : **>100,000 colonies** of one species
- ▶ Treatment can be delayed until culture results available.
- ▶ Positive culture (bacteriuria) alone **not** a reason to treat.

SPICE 

26

## McGeer Criteria


- ▶ Must fulfill both 1 AND 2
  - ▶ 1. At least 1 of the following signs/symptoms
    - ▶ Acute dysuria or pain, swelling, or tenderness of testes, epididymis, or prostate
    - ▶ Fever or leukocytosis and  $\geq 1$  of the following:
      - ▶ • Acute costovertebral angle pain or tenderness
      - ▶ • Suprapubic pain
      - ▶ • Gross hematuria
      - ▶ • New or marked increase in incontinence
      - ▶ • New or marked increase in urgency
      - ▶ • New or marked increase in frequency
    - ▶ If no fever or leukocytosis, then  $\geq 2$  of the following:
      - ▶ • Suprapubic pain • Gross hematuria • New or marked increase in incontinence • New or marked increase in urgency • New or marked increase in frequency
  - ▶ 2. At least 1 of the following microbiological criteria:
    - ▶  $\geq 10^5$  cfu/mL of no more than 2 species of organisms in a voided urine sample
    - ▶  $\geq 10^2$  cfu/mL of any organism(s) in a specimen collected by an in-and-out catheter



27

## Treatment /NO Treatment

- Asymptomatic bacteriuria should **NOT** be treated.
- Routine or post-treatment screening for bacteriuria is not recommended. (Infectious Diseases Society of America)
- ▶ No benefits in decreasing rates of subsequent UTIs
- ▶ Increased risk of resistance and uropathogens



28

## Indwelling Catheter-Associated UTI (CAUTI)


- ▶ Catheter colonization and infection is inevitable and expected!
- ▶ Once bacteria colonizes urine, concentration is 100,000 colonies within 72 hours!!



29

## Mechanisms of Colonization

- ▶ **Colonic and perineal flora primary source**
- ▶ Extra-luminal-- women – shorter urethra
- ▶ Manipulation of the collection system
- ▶ From hands of personnel during insertion
- ▶ Ascending from drainage bag/junction



30

## CMS UTI Antibiotic Treatment

Minimum criteria for initiating antibiotics for UTI

### **NO indwelling catheter, include:**

1. acute dysuria alone or fever ( $>37.9^{\circ}\text{C}$  [ $100^{\circ}\text{F}$ ] or  $1.5^{\circ}\text{C}$  [ $2.4^{\circ}\text{F}$ ] increase above baseline temperature) and at least one of the following:
  - new or worsening urgency, frequency, suprapubic pain, gross hematuria, costovertebral angle tenderness, or urinary incontinence.

Reference - "Development of Minimum Criteria for the Initiation of Antibiotics in Residents of Long-Term-Care Facilities: Results of a Consensus Conference" - *Infect Control Hosp Epidemiol* 2001;22:120-124.



31

## CMS UTI Antibiotic Treatment

Minimum criteria for initiating antibiotics for UTI

**2. Chronic indwelling catheter** (indwelling Foley catheter or a suprapubic catheter), includes the presence of at least one of the following:


- ▶ fever ( $>37.9^{\circ}\text{C}$  [ $100^{\circ}\text{F}$ ] or  $1.5^{\circ}\text{C}$  [ $2.4^{\circ}\text{F}$ ] increase above baseline temperature),
- ▶ new costovertebral tenderness, rigors (shaking chills) with or without identified cause, or new onset of delirium."

▶ Reference - "Development of Minimum Criteria for the Initiation of Antibiotics in Residents of Long-Term-Care Facilities: Results of a Consensus Conference" - *Infect Control Hosp Epidemiol* 2001;22:120-124.



32

## Intermittent Catheterization



- ▶ Intermittent catheterization can often manage overflow incontinence effectively.
  - ▶ New onset incontinence from a transient, hypotonic/atonic bladder (usually seen following indwelling catheterization in the hospital) may benefit from intermittent bladder catheterization until the bladder tone returns (e.g., up to approximately 7 days).
  - ▶ A voiding trial and post void residual can help identify when bladder tone has returned.
- overflow incontinence effectively



33

## USE of Urinary Catheters

### APPROPRIATE

- ▶ Clinical criteria for long/short for indwelling catheter:
  - ▶ Obstruction
  - ▶ Neurogenic bladder
  - ▶ Hematuria (short term)
  - ▶ Wounds stage 3 or >
  - ▶ Aggressive diuresis / monitoring of strict I/O (short term)
  - ▶ Terminally ill for comfort measures

### INAPPROPRIATE

- ▶ Convenience
- ▶ Used in lieu of other bladder management strategies.
- ▶ Used for specimen collection when the resident can voluntarily void

(Indwelling catheters are associated with a 5% risk/day of new UTI)



34

## CDC Guidelines on Flushing and irrigation

- ▶ If obstruction or infection occurs - change the catheter.

*"Unless obstruction is anticipated (e.g., as might occur with bleeding after prostatic or bladder surgery) bladder irrigation is not recommended...If obstruction is anticipated, closed continuous irrigation is suggested to prevent obstruction."*

- ▶ **"Q2C.3. Bladder irrigation**  
Low-quality evidence suggested no benefit of bladder irrigation in patients with indwelling or intermittent catheters.

[www.cdc.gov/infectioncontrol/pdf/guidelines/...](http://www.cdc.gov/infectioncontrol/pdf/guidelines/...)



35

## Prophylaxis For UTI Prevention: Cranberry

- ▶ **Cranberry juice/extract –**
  - ▶ Recent systematic reviews, including a 2023 update from the [Cochrane Database](#), suggest that cranberry products can help prevent urinary tract infections (UTIs) in certain populations.
  - ▶ The active ingredients, proanthocyanidins (PACs), work by preventing *E. coli* bacteria from adhering to the bladder wall
- ▶ **Women with recurrent UTIs:** Multiple studies indicate that cranberry products likely reduce the risk of UTIs in women with a history of frequent infections. A 2023 Cochrane review found a 26% reduced risk in this group.
- ▶ **Elderly or institutionalized patients:** The evidence does not support the use of cranberry products for UTI prevention in other populations such as elderly or institutionalized patients. Cranberry products may provide little to no benefit for older people living in nursing or care homes.



Williams G, Stothart CI, Hahn D, Stephens JH, Craig JC, Hodson EM. Cranberries for preventing urinary tract infections. Cochrane Database of Systematic Reviews 2023, Issue 11. Art. No.: CD001321. DOI: 10.1002/14651858.CD001321.pub7.



36

## Prophylaxis for UTI Prevention: Estrogens

- ▶ **Oral Estrogens** not shown to be beneficial.
- ▶ Topical, **vaginally applied estrogens** have been shown to be effective – 6 studies applying estrogen by ring, cream, or intravaginal tablet
  - ▶ Restores healthy vaginal tissue : keeps the tissue of the bladder and urethra healthy and flexible, improving function
  - ▶ Lowers vaginal pH which promotes growth of beneficial bacteria (like lactobacilli) and discourages pathogen growth



37

## Prophylaxis for UTI prevention: Methenamine

- ▶ A non-antibiotic medication used to prevent UTIs
  - ▶ Works by releasing formaldehyde in acidic urine (pH 5.5 or less)
  - ▶ Bc mechanism is different than antibiotics so bacteria don't develop a resistance
- ▶ Can lead to formation of uric acid crystals: Kidney stones: Lots of water necessary!
- ▶ Side effects: nausea, stomach upset, cramping
- ▶ Not effective for folks with renal tract abnormalities (neurogenic bladder)
- ▶ Methenamine vs Antibiotics in NH Patients (ALTAR Trial) : Effective in older adults! Even with reduced GFR.

Saul H, et al. C. Methenamine is as effective as antibiotics at preventing urinary tract infections. BMJ. 2023 Jan 17;380:72.  
 Harding C, et al. Alternative to prophylactic antibiotics for the treatment of recurrent urinary tract infections in women: multicentre, open label, randomised, non-inferiority trial. BMJ 2022;376:e068229.



38

## Prevention of UTI or Overtreatment

- ▶ Risk factor: Colonization
- ▶ Risk factor: Yeast
- ▶ Risk factor: Vaginal atrophy
- ▶ Risk factor: Indwelling Catheter
- ▶ Risk factor: Poor hygiene
- ▶ Prevention: Documentation
- ▶ Prevention: Await cultures
- ▶ Treatment: Vaginal estrogen, Vaseline
- ▶ Prevention: Remove catheter
- ▶ Prevention: Peri care and staff hand hygiene



39


## De-escalation in Urinary Tract Infection

1. Shorter length of therapy
  - Standard of care depends on the antibiotic choice, but is now typically 3 or 5 days.
  - Minimum necessary is best
2. Narrowing of spectrum
  - Utilize the culture results.
  - Consider awaiting treatment until these culture results return to ensure the appropriate antibiotic is being utilized.
3. Is this truly a UTI?

Goebel MC, Trautner BW, Grigoryan L. The Five Ds of Outpatient Antibiotic Stewardship for Urinary Tract Infections. Clin Microbiol Rev. 2021 Dec 15;34(4):e0000320. doi: 10.1128/CMR.00003-20. Epub 2021 Aug 25. PMID: 34431702; PMCID: PMC8404614.



40



41

### A Common Case

- ▶ 84 yo F living in your facility is “sleepier” today per son’s report. Staff note she has eaten less than usual and does seem more fatigued but has no other symptoms or complaints.
- ▶ Son insists that this is what happens “every time Mom has a urine infection” and requests that antibiotic treatment be initiated now. How do you respond?

SPICE

42

## Questions and Discussion

