

PRINCIPLES OF ANTIBIOTIC USE

REBEKAH MOEHRING, MD, MPH; DUKE UNIVERSITY MEDICAL CENTER
MEDICAL DIRECTOR, ANTIMICROBIAL STEWARDSHIP AND EVALUATION TEAM
SPICE 2026



dcasip.medicine.duke.edu



1

Disclosures

Grants to Institution: CDC

Royalties: UpToDate, Inc.



2

Objectives

Understand why antibiotics are “special” medications

The 4 Moments of Antibiotic Decision-Making

Antimicrobial Stewardship Programs

“Action” strategies in different clinical settings



3

We Love Antibiotics

Inpatient

At any given time, 60% of inpatients at DUH are receiving at least one antimicrobial

There are >26,000 antimicrobial admissions at DUH annually

30% of inpatient antimicrobial prescriptions are suboptimal

Long-term Care

Up to 50% of residents in a nursing home receive one or more courses of systemic antibiotics when followed over a year

40-75% of antibiotic prescriptions are inappropriate

Outpatient

423-553 antibiotic prescriptions per 1000 people in the US per year

28% are unnecessary, (representing 47 million prescriptions/year)



4

Why We Love Antibiotics

Wonder Drug

Active intervention

Experiences

Tangible

Insurance



Antibiotics are time-tested
placebos

Antibiotic Rx is easy:

- Avoids doing a structured exam or long DDx
- Avoids time-consuming discussions
- i.e. Easier to treat than diagnose or educate

Identifying Infected vs. Not
Infected is hard

“Just in case” perceived to be
lower risk than “watchful waiting”

5

Why we HAVE TO improve Antibiotic Use

Antibiotics are unlike any other drug, in that the use of the agent in one patient can compromise its efficacy in another.

A lot of antibiotic prescriptions are unnecessary or sub-optimal.

Antibiotic misuse harms patients.

Antibiotic misuse harms *future* patients.

Improving antibiotic use has many benefits for patients and society.



6

Antimicrobial Use Impacts: Infection Prevention, HAIs, AND Patient Outcomes



- Drug-resistance (MRSA, VRE, CRE, MDR-GNs)
- C. difficile* infection
- Infection treatment success/failure
 - Complications
 - Readmissions
 - Mortality
 - Length of Stay
- Adverse Safety Events
 - Allergic reactions
 - Drug toxicity events
 - Acute Kidney Injury
- Healthcare Resources and Cost
 - (all of the above)
 - Pharmacy budget; ICU days



7


One in Five Inpatients get an Antibiotic Adverse Drug Event

1488 patients followed for 30 days after antibiotic initiation

Followed 90 days for CDI and MDRO acquisition

General medical inpatients who had at least 24h of antibiotics during admission


20% of patients experienced at least one antibiotic-associated ADE



Tamma et al. JAMA Int Med. 2017 177(9):1308-15.



8



9

A case

86F with history of dementia, diabetes, and poor functional status presents from SNF with confusion, fever.

ED: Hypotense, non-verbal. Fluid resuscitated, cultures drawn, started on vancomycin + zosyn and admitted to the floor with diagnosis of sepsis.

Day 3: Remains on vancomycin and zosyn, progress note still says "sepsis." BCx negative. Awake/interactive.

Urine culture: E. coli susceptible to multiple oral and intravenous agents.

10

Making the Right Decision Is Important


Getting the correct antibiotic to our sickest patients in a timely manner is critical.

Inappropriate empiric antibiotics associated with increased mortality and longer length of stay.

- In febrile inpatients, in ICU patients

In Septic Shock, each hour delay in administration of appropriate antimicrobials is associated with increased mortality.

- Average decrease in survival of 7.6%/hour over the first 6h after shock onset



Surviving Sepsis Campaign

Fraser A, *et al.* Am J Med 2006;119:970-6
 Kollef, M, *et al.* Chest 1999. 115: 462-474
 Kumar *et al.* Crit Care Med. 2006 Jun;34(6):1589-96.

11

+You Search Images Maps Play YouTube Gmail Drive More ▾

Google sepsis antibiotics

Web Images Videos News Shopping Maps Books

About 69,564,000 results (0,81 seconds)

Did you mean: **vanc/zosyn**

Created with didyoumean-generator.com

Duke Center for Antimicrobial Stewardship and Infection Prevention

12

But, not for ALL infected patients

“Sepsis” without shock – a very heterogeneous population that does NOT show the same time-related associations with antibiotic/mortality

Sepsis can be hard to diagnose. Many patients with hypotension have non-infectious diagnoses.

- pulmonary emboli, fluid under/overload, toxin exposures, drug adverse effects, malignancies, bleeding, mechanical complications of surgery, obstructed organs, etc....

Overly tight limits on timing of antibiotics for suspected sepsis may cause clinicians to err on the side of over-treatment, skip diagnostic steps, and subject patients to the harms of antibiotic overuse.

Weinberger et al. J Infect Dis. 2020 Jul 21;222(Suppl 2):S110-S118
Liu VX. Am J Respir Crit Care Med 2017; 196:856–863
Seymour CW, et al. N Engl J Med 2017; 376:2235–2244



13

... And not Broadest Spectrum Abx for All

“Suggest against” Empiric Coverage of MDR Pathogens unless deemed “High risk”:

- Personal prior history (e.g. known prior positive culture with PsA)
- Known colonization with MDRO (e.g. Nares PCR for MRSA+)
- Prior prolonged use of broad-spectrum antibiotics
- Prior prolonged stay in hospital environment with high prevalence of MDRO (e.g. >20%)

Empiric Multidrug Resistant (MDR) Pathogen Coverage

-

25. For adults with sepsis or septic shock at high risk of infection with a specific multidrug resistant (MDR) pathogen, we “suggest” using empirical antimicrobial therapy with coverage for this MDR pathogen.
(conditional recommendation, very low certainty evidence)

Remark: Risk factors for MDR pathogens include colonization with the MDR pathogen of concern, previous infection with the MDR pathogen of concern, prolonged use of broad-spectrum antibiotics, and prolonged hospitalization in a unit with a high prevalence of the MDR pathogen of concern.

26. For adults with sepsis or septic shock at low risk of infection with a specific MDR pathogen, we “suggest against” using empirical antimicrobial therapy with coverage for this MDR pathogen.
(conditional recommendation, very low certainty evidence)

Surviving Sepsis Guidelines 2026



14


Undifferentiated, Suspected Sepsis

- Cross sectional study of **6.3 million adults across 241 US hospitals** between 2017-2021
- Focused on **empiric choices** of antibiotics
- **Half** of all anti-methicillin resistant *Staphylococcus aureus* and antipseudomonal B-Lactam antibiotics were **prescribed for suspected community onset sepsis**
- **Only 9.5%** of suspected community onset sepsis treated with broad spectrum antibiotics **had resistant organisms identified**

Rhee et al. JAMA NO 2024. (7):6



Figure 4. Trends in Empiric Broad-Spectrum Antibiotic Use for Community-Onset Suspected Sepsis.



[Open in a new tab](#)

MRSA methicillin-resistant *Staphylococcus aureus*.

15

AU represents a modifiable risk

AU in Nursing Homes is highly variable and correlated with AEs

Figure. Variability of Antibiotic Use (per 1000 Resident-days) Across Ontario Nursing Homes




Table 3. Antibiotic-Related Adverse Outcomes Among Residents Living in Nursing Homes With Low, Medium, and High Antibiotic Use^a


Characteristic	Antibiotic Use, No. (%)		
	Low (n = 33 822)	Medium (n = 31 425)	High (n = 24 943)
<i>Clostridium difficile</i>	274 (0.8)	268 (0.9)	221 (0.9)
Diarrhea or gastroenteritis	3347 (9.9)	3388 (10.8)	2889 (11.6)
Infection with antibiotic-resistant organism	412 (1.2)	431 (1.4)	319 (1.3)
Antibiotic allergy	13 (0.0)	25 (0.1)	22 (0.1)
General adverse event from medication	96 (0.3)	124 (0.4)	88 (0.4)
Any antibiotic complication with or without potential for indirect harms to nonrecipients (primary composite outcome ^b)	3869 (11.4)	3890 (12.4)	3311 (13.3)
Only antibiotic complications with potential for indirect harms to nonrecipients (secondary composite outcome ^c)	3797 (11.2)	3801 (12.1)	3237 (13.0)

Daneman et al. JAMA IM 2015;175 (8): 1331-1339



16

Acute Care Academic Hospitals



17

Good use of antibiotics requires balance. It's nuanced and complex.

Right Diagnosis

Right Drug


Right Dose

Right Timing

Right Duration


Improve therapeutic choices (underuse)

Reduce unnecessary use (overuse)



18

The “4 Moments” of Antibiotic Decision-Making



Tamma PD et al. JAMA. 2019;321(2):139-140.

5 or 6 “Ds” of Antimicrobial Stewardship

Morency-Potvin, Schwartz, Weinstein. Clin Micro Rev 30: 381-407

Diagnosis	Make and document the right diagnosis.
Debridement/ Drainage	Drainage of abscesses and removal of necrotic tissue or foreign material when required.
Drug	Use the right drug empirically according to suspected or confirmed diagnosis, risk factors for resistant pathogens, allergy, or major side effects.
Dose	Use right dose according to diagnosis, site of infection, or renal/hepatic dysfunction.
Duration	Use drugs for an appropriate duration.
De-escalation	Re-evaluate diagnosis and therapy routinely and de-escalate therapy to narrow-spectrum and/or oral agents when appropriate.

Goebel et al. Clin Micro Rev 2021. <https://doi.org/10.1128/CMR.00003-20>

Diagnosis	Make and document the right diagnosis
Drug	Use the right empiric antibiotic
Dose	Use the right dose of antibiotic based on site of infection and renal or hepatic dysfunction
Duration	Use antibiotics for the recommended duration
De-escalation	De-escalate therapy based on susceptibilities and when urine cultures are negative



Clinical information: Small pieces over time.

Clinical information trickles in over time.

This means clinicians have to reassess regularly.

This also means they get interrupted with 'real-time' notifications and need to respond.

This a complex process: unpredictable, unknowns, uncertainty.

Putting the puzzle together completely takes attention, follow up on details, ability to make decisions in the setting of unknowns, AND an eye on the long-term goals.



21

General Indications for Antibiotics

Prophylaxis: prevent infection

- EASY! Guidelines and ordersets

Empiric: when you suspect infection but don't exactly know with what pathogen

- Not easy. Local guidelines help (based on local micro data).

Directed: pathogen known

- Moderately easy. Follow and interpret patient-specific micro data.

2

3


Have I ordered appropriate cultures before starting antibiotics?

What empirical antibiotic therapy should I initiate?

A day or more has passed.

Can I stop antibiotics?
Can I narrow therapy?
Can I change from IV to oral therapy?

Empiric
Broad-spectrum



Targeted
Narrow-spectrum



22

Choice of Empiric Antimicrobials: Clinical Syndrome

What class of pathogen am I likely to be treating?

- (Bacterial? Viral? Fungal? Other?)

2

If bacterial, what organisms are most likely?

- (Gram positive? Gram negative? Anaerobe?)

What information can I get to guide treatment?

- Microbiology data?

Do I need to order any other diagnostic tests?

How sick is my patient? How risky would it be if I miss?

Is my patient “special”? – allergy, ADEs, immune status

Have I ordered appropriate cultures before starting antibiotics?

What empirical antibiotic therapy should I initiate?




23

Traditional Microbiologic Culture

- “Gold standard” for diagnosis
- Requires sampling of site of infection, ideally *prior to* therapy
- Allows determination of phenotypic antimicrobial susceptibility

Growth? Up to 5 days



Rapid
PCR-
based
Test!

ID most common pathogens
and *Genotypic* resistance



24

DIAGNOSIS: Stain

Direct Visualization

Gram stain

- Often provide clues to etiology (may allow presumptive diagnosis in some cases)

- Gram Positive
- Gram Negative
- Non-staining

Shape


- Cocci
- Rods

Aerobic/Anaerobic

Disclaimer: We use Gram Stain in our *conceptual* understanding of Pathogens/Abx Coverage...

BUT

Rapid molecular diagnostic tests provide quicker info about common pathogen IDs and genotypic resistance



www.laboratoryinfo.com

25

Quick and Dirty Anti-bacterial Classification

Gram positive – skin, lung, guts, devices

Gram negative – guts, urine, some lung, chronic wounds

Atypicals – lung, STIs

Anaerobes –gas- and abscess-forming, bad odors

Fungi – guts, devices, immunosuppressed + abx-exposed hosts



26

Pathogens are tricky.

Antibiotic = A drug that kills or inhibits the growth of bacterial pathogens

Resistant = Designation that implies that an antimicrobial will not inhibit bacterial growth at clinically achievable concentrations

Susceptible = Designation that implies that an antimicrobial will inhibit bacterial growth at clinically achievable concentrations

Susceptibility

	Serratia marcescens MIC
Amikacin	S
Ampicillin	R
Ampicillin + Sulbactam	R
Cefazolin	R
Ceftriaxone	S
Ciprofloxacin	I
Gentamicin	S
Piperacillin/Tazobactam	S
Tobramycin	S
Trimethoprim + Sulfamethoxazole	S



27

Patients are individuals.

Drug interactions

Age

Allergies

Pregnancy, breast feeding

Toxicity (idiosyncratic reactions)

Dose adjustment for renal and/or hepatic dysfunction

Ability to absorb an oral antibiotic

Immune status

Adherence:

- Cost
- Taste
- Frequency of administration
- Pill size
- Duration of therapy
- Multiple drug therapy
- Adverse effects
- Current symptoms



28

De-escalation

De-escalation is a core principle of Antimicrobial Stewardship.

Target/narrow antibiotic therapies after more clinical data returns

Stop therapy when infection has been ruled out

Goal: Reduce selective pressure, improve safety



29

Durations (OLD)

- Most guideline-recommended antibiotic durations are based on...
- Qualified with “it depends...”
- Duration questions are ~70% of ID consults.

The answer, Dr. Gilbert, is that this is highly-specialized knowledge, rarefied information that only 100% Board-Certified, USDA-inspected Infectious Diseases Doctors know. And since I'm concerned that your article might give readers the wrong impression about our scientific credibility, I'll now divulge what we've learned, and how to apply it.

To figure out how long antibiotics need to be given, use the following rules:

1. Choose a multiple of 5 (fingers of the hand) or 7 (days of the week).
2. Is it an outpatient problem that is relatively mild? If so, choose something less than 10 days. After application of our multiples rule, this should be 5 or 7 days.
3. Is it *really* mild, so much so that antibiotics probably aren't needed at all but clinician or patient are insistent? Break the 5/7 rule and go with 3 days. Ditto uncomplicated cystitis in young women.
4. Is it a serious problem that occurs in the hospital or could end up leading to hospitalization? With the exception of community-acquired pneumonia (5 or 7 days), 10 days is the minimum.
5. Patient not doing better at the end of some course of therapy? Extend treatment, again using a multiple of 5 or 7 days.
6. Does the infection involve a bone or a heart valve? Four weeks (28 days) at least, often 6 weeks (42 days). Note that 5 weeks (35 days) is not an option — here the 5's and 7's cancel each other out, and chaos ensues.
7. The following lengths of therapy are inherently weird, and should generally be avoided: 2, 4, 6, 8, 9, 11, 12, 13 days. Also, 3.14159265 days.

In this highly data-driven exercise, it is important also to note the *number* of rules — *seven*, as in days of the week.

<https://blogs.jwatch.org/hiv-id-observations/index.php/how-to-figure-out-the-length-of-antibiotic-therapy/2010/10/22/>



Paul E. Sax, MD

Contributing Editor
NEJM JOURNAL WATCH
INFECTIOUS DISEASES



30

NOW: Trials comparing short- vs. longer-course antibiotics have shown short-course is just as effective

4

What duration of antibiotic therapy is needed for this patient's diagnosis?

Disease	Antibiotic Duration	
	Short	Long
Community-acquired pneumonia	3-5 days	7-10 days
Nosocomial pneumonia (HAP/VAP)	≤ 8 days	10-15 days
Pyelonephritis	5-7 days	10-14 days
Intraabdominal infection	4 days	10 days
Acute exacerbation of chronic bronchitis (AECB) and COPD	≤ 5 days	≥ 7 days
Acute bacterial sinusitis	5 days	10 days
Cellulitis	5-6 days	10 days
Chronic osteomyelitis	42 days	84 days
Gram-negative bacteremia	7 days	10-14 days



Spellberg B. JAMA Intern Med 2016;176(9):1254-1255.

31

WHAT IS ANTIMICROBIAL STEWARDSHIP?



dcasip.medicine.duke.edu



32

IDSA/SHEA/PIDS definition

“coordinated interventions designed to improve and measure the appropriate use of [antibiotic] agents by promoting the selection of the optimal [antibiotic] drug regimen including dosing, duration of therapy, and route of administration.”

-- *Infectious Diseases Society of America (IDSA), the Society for Healthcare Epidemiology of America (SHEA), and the Pediatric Infectious Diseases Society (PIDS)*

Barlam et al. *CID* 2016; 62(10): e51-77.
Fishman et al. *ICHE* 2012; 33:322-7.



33

Antimicrobial Stewardship Program

Support system for prescribers of antimicrobials.

Coordinated program

Multidisciplinary teams

- MD, PharmD, RN, micro, IP, IT

Multi-level interventions:

- Educational
- Systems-based vs. 1:1
- Technology
- Active vs. Passive



Dellit et al. *Clin Inf Dis.* 2007;44(2):159-177.



34

US Hospitals meeting CDC's 7 Core Elements of AS

NC: 100% in 2024

<https://arpsp.cdc.gov/profile/stewardship>

Percentage of hospitals meeting All 7 Core Elements

Year

37

CDC 2022 "Priorities for Core Elements Implementation"

<https://www.cdc.gov/antibiotic-use/core-elements/hospital/priorities.html>


Table. The Core Elements of Hospital Antibiotic Stewardship Programs and the Priorities for Hospital Core Element Implementation.

Hospital Core Elements	Priorities for Hospital Core Element Implementation
Hospital Leadership Commitment Dedicate necessary human, financial, and information technology resources.	Antibiotic stewardship physician and/or pharmacist leader(s) have antibiotic stewardship responsibilities in their contract, job description, or performance review.
Accountability Appoint a leader or co-leaders, such as a physician and pharmacist, responsible for program management and outcomes.	Antibiotic stewardship program is co-led by a physician and pharmacist.*
Pharmacy/Stewardship Expertise Appoint a pharmacist, ideally as the co-leader of the stewardship program, to help lead implementation efforts to improve antibiotic use.	Antibiotic stewardship physician and/or pharmacist leader(s) have completed infectious diseases specialty training, a certificate program, or other training on antibiotic stewardship.
Action Implement interventions, such as prospective audit and feedback or preauthorization, to improve antibiotic use.	Antibiotic stewardship program has facility-specific treatment recommendations for common clinical condition(s) and performs prospective audit/feedback or preauthorization.
Tracking Monitor antibiotic prescribing, impact of interventions, and other important outcomes, like C. difficile infections and resistance patterns.	Hospital submits antibiotic use data to the NHSN Antimicrobial Use Option.
Reporting Regularly report information on antibiotic use and resistance to prescribers, pharmacists, nurses, and hospital leadership.	Antibiotic use reports are provided at least annually to target feedback to prescribers. In addition, the antibiotic stewardship program monitors adherence to facility-specific treatment recommendations for at least one common clinical condition.
Education Educate prescribers, pharmacists, nurses, and patients about adverse reactions from antibiotics, antibiotic resistance, and optimal prescribing.	No implementation priority identified.

* For critical access hospitals (CAHs), this criterion can be met if the hospital has a physician leader with a pharmacist involved in stewardship (recognizing that some CAHs do not have pharmacists on staff, so co-leadership is not possible).

1. Dedicated FTE AS in contract/performance evaluation
2. Accountable leaders: physician and pharmacist
3. Expertise: in ID, or certificate
4. Do something: "Core" strategies (PA or PAF) + Local GL
5. Track: AU Rates (via NHSN) + resistance + HAIs
6. Share data back: annual, monitor adherence to local GL
7. Do some education

38



39

Priority “Action” Strategies for Inpatient Stewardship

Preauthorization/restriction (before prescribing)

- Can be time/personnel intensive
- Must think through unintended consequences and process snafus
- Not for all hospitals – local culture plays a role
- Best if for targeted agents (not every antibiotic order...)

Post-prescription audit and feedback (after prescribing)

- Front-line stewardship “experts” actively review patients on antibiotics and give feedback to prescribers 1:1
- Time intensive, but effective
- Better for personal relationships
- Need ID consult “back up” for tough cases

**“Handshake”
Stewardship
or
Stewardship Rounds**

Face to face is better than
phone/pager/zoom.

CID 2016;62(10):e51–e77
 Hurst A et al. PIDJ 2016;35(10):1104-10
 Seidelman. CID 2022; 74(11):1986-1992




40

Monthly Prospective Reviews:

Stewardship Interventions (Adult and Pediatrics)



*% of reviewed patients had a least 1 intervention.



1. Antibiotic streamlining/de-escalation (35%)
2. Dose/Drug (25%)
3. Duration/Discontinuation (17%)
4. Allergy assessment (9%)
5. ID consult rec (5%)
6. IV/PO switch (3%)



41

REVISED TJC Inpatient Stewardship Standards: Jan 2023

MM.09.01.01



Standards related to measurement of AU:

- 15. The antibiotic stewardship program documents the evidence-based use of antibiotics in all departments and services of the hospital.
- 16. The antibiotic stewardship program monitors the hospital's antibiotic use by analyzing data on days of therapy per 1000 days present or 1000 patient days, or by reporting antibiotic use data to the National Healthcare Safety Network's Antimicrobial Use Option of the Antimicrobial Use and Resistance Module.
- ...
- 19. The antibiotic stewardship program evaluates adherence (including antibiotic selection and duration of therapy, where applicable) to at least one of the evidence-based guidelines the hospital implements.
 - Note 1: The hospital may measure adherence at the group level (that is, departmental, unit, clinician subgroup) or at the individual prescriber level.
 - Note 2: The hospital may obtain adherence data for a sample of patients from relevant clinical areas by analyzing electronic health records or by conducting chart reviews.

42

Local Guidelines + Adherence Reporting Priority for Inpatient Stewardship

Facility-specific treatment recommendations for common clinical condition(s)

AMERICAN THORACIC SOCIETY DOCUMENTS

Diagnosis and Treatment of Adults with Community-acquired Pneumonia
An Official Clinical Practice Guideline of the American Thoracic Society and Infectious Diseases Society of America

Joshua P. Metlay*, Grant W. Waterer*, Ann C. Long, Antonio Anzueto, Jan Brozek, Kristina Crothers, Laura A. Cooley, Nathan C. Dean, Michael J. Fine, Scott A. Flanders, Marie R. Griffin, Mark L. Metersky, Daniel M. Musher, Marcos I. Restrepo, and Cynthia G. Whitney, on behalf of the American Thoracic Society and Infectious Diseases Society of America

THIS OFFICIAL CLINICAL PRACTICE GUIDELINE WAS APPROVED BY THE AMERICAN THORACIC SOCIETY MAY 2019 AND THE INFECTIOUS DISEASES SOCIETY OF AMERICA AUGUST 2019



Local Patient Population(s)
Local Formulary
Local antibiogram

Typically requires pharmacy + physician input, approved by hospital or system committee

Dissemination
Education
Integration into ordersets/pathways

****Assessment of Adherence (2023)**

43

<https://www.cdc.gov/nhsn/PDFs/pscManual/11pscAURcurrent.pdf>


“Tracking” Antibiotic Use through NHSN

NHSN AU Option for acute care hospitals

- ONLY uses electronic data from EHRs (no manual surveillance and no subjective components)

Rate: Days of therapy (DOT) per 1,000 days present

- DOT = calendar days of treatment regardless of number of doses
- Separate drugs counted separately
- Denominator is DIFFERENT than patient days



Data is stratified by Agent, Route, Unit location

Benchmark: Standardized Antimicrobial Administration Ratio (SAAR)



44

Benchmarking AU between Hospitals

What you want to measure:

- Prescribing practices and decision-making


What you get:

- Tonnage or measure of abx exposure – not “appropriate” abx

Problems:

- Case mix
- Hospital size
- Clinical service lines (e.g. surgical specialties, types of ICUs, moms/babies)
- Assume more = bad (not always true clinically)

Should be viewed as a starting point for further investigation and explanation.



Ibrahim, Polk. Expert Rev Anti Infect Ther 10 (4):445-57. Stenehjem. CID 2016;63(10):1273-80

45

NHSN Standardized Antibiotic Administration Ratio (SAAR)

Standardized Antibiotic Administration Ratio: Observed/Expected based on NHSN baseline + limited risk adjustment with information from annual survey

- Heavily stratified: Pediatric (8), NICU (7), and Adult (7) agent groups; Pediatric (5), NICU (3), and Adult (8) NHSN unit types. *New benchmarks for more specialized units released this year.

Baseline 2017 – RECENT REBASELINE to 2023**

- Most current SAAR Targets are pooled mean + percentiles reported in annual report (2024)

<https://www.cdc.gov/nhsn/datastat/aur-reports.html>

<https://www.cdc.gov/nhsn/pdfs/ps-analysis-resources/aur/au-saar-guide-508.pdf>

Broad spectrum antibacterial agents predominantly used for hospital-onset infections used in adult SAAR wards

Facility Org ID	SAAR Type 2017 Baseline	Location	Summary Year/Month	CDC Location	Antimicrobial Days	Predicted Antimicrobial Days	Days Present	SAAR	SAAR p-value	95% Confidence Interval	SAAR Percentile
10009	Adult_BSHO_Ward_2017	MEDSURG64	2022M01	IN:ACUTE:WARD:MS	45	36.819	320	1.222	0.1852	0.902, 1.621	64



46

Antibiotic Use And Resistance Reporting Is Required for Acute Care Hospitals via CMS

The requirement was in the 2023 Inpatient Prospective Payment System rule from CMS. The requirement is under the Promoting Interoperability program standard “Public Health and Clinical Data Exchange Objective”

<https://www.federalregister.gov/documents/2022/08/10/2022-16472/medicare-program-hospital-inpatient-prospective-payment-systems-for-acute-care-hospitals-and-the#h-623>

Not only AU Option, but also the AR module

➤ Hospitals that don't participate lose their incentive (\$\$\$) by Jan 2025 (for reporting year 2024).

Page 49335
Slide: Arjun Srinivasan



Measuring “Appropriate” or “Optimal” antimicrobial use in the Inpatient Setting

Will generally introduce subjectivity and require individual case review by a trained individual

More focused to syndrome and/or drug

- “MUE” + adherence with local guidelines

Typically sampled prevalence or retrospective study, but can be prospective and integrated into intervention delivery

Big time personnel effort

		If endorsed guidelines are present	If endorsed guidelines are absent
Appropriate	1 Optimal ¹	Antimicrobial prescription follows either the Therapeutic Guidelines ² or endorsed local guidelines optimally, including antimicrobial choice, dosage, route and duration ³	The antimicrobial prescription has been reviewed and endorsed by an infectious diseases clinician or a clinical microbiologist OR The prescribed antimicrobial will cover the likely causative or cultured pathogens and there is not a narrower spectrum or more appropriate antimicrobial choice, dosage, route or duration ³ available
	2 Adequate	Antimicrobial prescription does not optimally follow the Therapeutic Guidelines ² or endorsed local guidelines, including antimicrobial choice, dosage, route or duration ³ ; however, is a reasonable alternative choice for the likely causative or cultured pathogens OR For surgical prophylaxis, as above and duration ³ is less than 24 hours	Antimicrobial prescription including antimicrobial choice, dosage, route and duration ³ is not the most optimal, however, is a reasonable alternative choice for the likely causative or cultured pathogens OR For surgical prophylaxis, as above and duration ³ is less than 24 hours
Inappropriate	3 Suboptimal	There may be a mild or non-life-threatening allergy mismatch OR Antimicrobial prescription including antimicrobial choice, dosage, route and duration ³ is an unreasonable choice for the likely causative or cultured pathogens, including: • spectrum excessively broad, unnecessary overlap in spectrum of activity, dosage excessively high or duration excessively long • failure to appropriately de-escalate with microbiological results	
	4 Inadequate	Antimicrobial prescription including antimicrobial choice, dosage, route or duration ³ is unlikely to treat the likely causative or cultured pathogens OR The documented or presumed indication does not require any antimicrobial treatment OR There may be a severe or possibly life-threatening allergy mismatch, or the potential risk of toxicity due to drug interaction OR For surgical prophylaxis, the duration ³ is greater than 24 hours (except where local guidelines endorse this)	
	5 Not assessable	The indication is not documented and unable to be determined from the notes OR The notes are not comprehensive enough to assess appropriateness OR The patient is too complex, due to multiple co-morbidities, allergies or microbiology results, etc.	

¹Taking into account acceptable changes due to the patient's weight, allergy status, renal or hepatic function, or relevant drug interactions (if this information is available)
²Antibiotic Expert Group, Therapeutic Guidelines: Antibiotic, Version 16 (2015), or online version
³Duration should only be assessed if the guidelines state a recommended duration and the antimicrobial has already been dispensed for longer than this, or if there is a clear planned 'end date' documented



<https://www.ncas-australia.org/ncas-publications>

CMS: ASP required in Long-term Care

CMS Requirement for Long-term Care ASPs since 2017

Barriers to Implementation of AS in LTC:

Knowledge/Evidence

Expertise

Different stakeholders + processes of care than acute care

Personnel and turnover



49

Examples of Stewardship “Action” in LTC

Antibiotic use protocols – “Minimum Criteria” for Abx starts

Test/diagnostic stewardship

- UA/culture
- *C. difficile*

Durations/length of therapy and guides for common infection

- UTI
- Pneumonia
- Cellulitis

“Active monitoring” as an alternative to empiric antibiotics in patients who have a clinically undifferentiated problem (e.g. “not at baseline”)



50

Example: Minimum Criteria Protocol

Target: Nursing assessment

Identifies “red flag” symptoms

Includes “Notes” that identify key areas for baseline knowledge

Provides next steps alternative (other than an antibiotic)

SBAR tool for communication to clinician

AHRQ Toolkit: “Minimum Criteria for Common Infections”



Minimum Criteria for Initiating Antibiotics for a Skin and Soft Tissue Infection

Initiate antibiotics if the following criteria are met:

- New or increasing purulent drainage at a wound, skin, or soft-tissue site

OR

- At least two of the following:
 - Fever (temperature > 100°F [37.9°C] or two repeated temperatures of 99°F [37°C]), or
 - Redness, or
 - Tenderness, or
 - Warmth, or
 - Swelling that is new or increasing at the affected site

Notes:

1. For residents that regularly run a lower temperature, use a temperature of 2°F (1°C) above the baseline as a definition of a fever.
2. Herpes zoster is a virus and therefore does not require antibiotics but appropriate antivirals.
3. Odor is not a standalone criterion for treatment with antibiotics
4. Deeper infections such as bursitis may present with similar signs/symptoms.
5. Underlying osteomyelitis should be considered when managing a resident with an infected diabetic or decubitus ulcer.
6. Thromboembolic disease should be considered when a resident presents with an erythematous or swollen leg.
7. These criteria do not apply to residents with burns.
8. Gout can at times be mistaken for cellulitis or vice versa.

If none of the minimum criteria are met, consider initiating the following:

- Assess vital signs, including temp, every ____ hours for ____ hours; and/or
- Notify Physician/NP/PA if symptoms worsen or if unresolved in ____ hours.

Regardless of whether the minimum criteria are met or not, consider initiating the following:

- For discomfort or prior to cleaning/dressing changes, consider using acetaminophen or other pain relievers as needed.

51

Example: Active Monitoring

This is an active process

More frequent vital signs

Oral hydration

Assess for pain, changes in medicine, other reasons like a bad night's sleep

(or disagreement with a loved one)




Order Set

- Obtain vital signs (BP, Pulse, Resp Rate, Temp, Pulse Ox) every ____ hours for ____ days.
- Record fluid intake each shift for ____ days.
- Notify physician if fluid intake is less than ____ cc daily.
- Offer resident ____ ounces of water / juice every ____ hours.
- Notify physician, NP, or PA if condition worsens, or if no improvement in ____ hours.
- Obtain the following blood work _____.
- Consult pharmacist to review medication regimen.
- Contact the physician, NP, PA with an update on the resident's condition on _____.

Nace *et al.*, JAMDA 2014;15:133-139

52




NIAID
Public Health Image
Library

Potential Policies & Procedures: UTI

- Concerns about stinky or cloudy urine should lead to increased hydration and perhaps, watchful waiting/careful observation.
- Automatic review of all medication changes by outside providers.
- Send residents to the Emergency Room with a note clearly stating what you are (and are not) worried about.

53



NIAID
Public Health Image
Library


Potential Policies & Procedures: UTI

- Clear criteria for collecting a urine sample
- Documented protocol for proper sample collection and handling
- Communication tools (SBAR) when nurses call a covering provider
- Proactively talk to residents and their family members—on admission and during change of status

54

Example: Education for Residents and Family


https://www.ahrq.gov/sites/default/files/wysiwyg/nhguide/6_TK1_T5-Suspect a Urinary Tract Infection brochure MA Coalition final.pdf



Developed by the Massachusetts Infection Prevention Partnership

Suspect a Urinary Tract Infection? How Taking Antibiotics When You Don't Need Them Can Cause More Harm Than Good


An Important Message for Seniors and their Families




Did You Know That...

- »Up to 50 percent of all antibiotics prescribed are not needed or are not prescribed appropriately?
- »Confusion or sudden behavior changes don't necessarily indicate a urinary tract infection (UTI)?
- »As many as half of seniors living in long-term care settings will test positive for bacteria in their urine, **without actually having a UTI?**

Learn Why The CDC is Sounding The Alarm
About The Overuse of Antibiotics



Massachusetts Coalition
for the
Prevention of Medical Errors



© Massachusetts Coalition for the Prevention of Medical Errors - used with permission of from the Coalition and the Massachusetts Infection Prevention Partnership

So if a test shows bacteria, does this mean I have a UTI?

Not necessarily. Diagnosis of a UTI requires **both** finding bacteria on a urine test **and** the presence of specific symptoms. Having **both** is important, because bacteria can and do live naturally in the bladder without causing any pain or symptoms. This is called **asymptomatic bacteriuria**, which is present in as many as half of seniors living in long-term care settings.

What are the specific symptoms of a UTI?

- » a burning feeling, discomfort or pain with urination.
- » pain in the lower abdomen or back.
- » increase in frequency (needing to "go" more often than usual).
- » repeated strong urges to urinate.
- » blood in the urine.

These symptoms may or may not be accompanied by fever.

What about other symptoms, such as confusion or a sudden change in behavior?

UTI is less likely without the specific symptoms listed above. Non-specific symptoms such as confusion, a sudden change in behavior, fatigue or a fall may be caused by other factors, including:

- » dehydration
- » inadequate nutrition
- » poor sleep
- » depression
- » medication side effects
- » constipation

It is important to consider a range of possible causes, to prevent missing the real diagnosis.

I was prescribed antibiotics before. Why not now?

In the past, when a urine specimen tested positive — *even when no symptoms of infection were present* — doctors were taught that treatment with antibiotics was the right approach. **We know now that is not correct.**

The American Geriatric Society now recommends to physicians
"Don't use antimicrobials (antibiotics) to treat bacteriuria in older adults unless specific urinary tract symptoms are present."

55

Resources for Outpatient Stewardship


Outpatient Standard for TJC (Jan 2020)

4 "Core Elements"

Type of outpatient practice setting is highly varied

- Adult/pediatric
- Specialty clinics
- Retail clinics
- Urgent Care

https://www.cdc.gov/antibiotic-use/community/pdfs/16_268900-A_CoreElementsOutpatient_508.pdf



Clinician Checklist for Core Elements of Outpatient Antibiotic Stewardship

CDC recommends that outpatient clinicians take steps to implement antibiotic stewardship activities. Use this checklist as a baseline assessment of policies and practices that are in place. Then use the checklist to review progress in expanding stewardship activities on a regular basis (e.g., annually).

COMMITMENT

1. Can you demonstrate dedication to and accountability for optimizing antibiotic prescribing and patient safety related to antibiotics? Yes No

If yes, indicate which of the following are in place (select all that apply)

- Write and display public commitments in support of antibiotic stewardship.

ACTION

2. Have you implemented at least one practice to improve antibiotic prescribing? Yes No

If yes, indicate which practices which you use. (Select all that apply.)

- Use evidence-based diagnostic criteria and treatment recommendations.
- Use delayed prescribing practices or watchful waiting, when appropriate.

TRACKING AND REPORTING

3. Do you monitor at least one aspect of antibiotic prescribing? Yes No

If yes, indicate which of the following are being tracked. (Select all that apply.)

- Self-evaluate antibiotic prescribing practices.
- Participate in continuing medical education and quality improvement activities to track and improve antibiotic prescribing.

EDUCATION AND EXPERTISE

4. Do you provide education to patients and seek out continuing education on antibiotic prescribing? Yes No

If yes, indicate how you provide antibiotic stewardship education. (Select all that apply.)

- Use effective communications strategies to educate patients about when antibiotics are and are not needed.
- Educate about the potential harms of antibiotic treatment.
- Provide patient education materials

56

“Action” in Outpatient Stewardship

Most literature in Primary or Urgent Care

Peer comparison + data feedback

- Most commonly done for upper respiratory infection
- Identify diagnoses (e.g. viral URI) for which antibiotics should not be given. Benchmark % given abx with peers
 - HEDIS measures (primary care and pediatrics)

Suggested alternatives

Accountable justification

“Nudge” letter/poster

Education combined with the above



Gerber. JAMA 2013;309(22):2345-2352
 Meeker. JAMA 2016; 315(6):562-570
 Meeker. JAMA Intern Med. 2014;174(3):425-431.

We want to give you some important information about antibiotics.

Antibiotics, like penicillin, fight infections due to bacteria that can cause some serious illnesses. But these medicines can cause side effects like skin rashes, diarrhea, or yeast infections. If your symptoms are from a virus and not from bacteria, you won't get better with an antibiotic, and you could still get these bad side effects.

Antibiotics also make bacteria more resistant to them. This can make future infections harder to treat. This means that antibiotics might not work when you really need them. Because of this, it is important that you only use an antibiotic when it is necessary to treat your illness.

How can you help? Carefully follow your doctor's instructions. He or she will tell you if you should or should not take antibiotics.

When you have a cough, sore throat, or other illness, your doctor will help you select the best possible treatments. If an antibiotic would do more harm than good, your doctor will explain this to you, and may offer other treatments that are better for you.

Your health is very important to us. As your doctors, we promise to treat your illness in the best way possible. We are also dedicated to avoid prescribing antibiotics when they are likely to do more harm than good.

If you have any questions, please feel free to ask your doctor, nurse, or pharmacist.

57

HOT in Stewardship RN



Diagnostic Stewardship/Diagnostic Excellence (DxEx)

- CDC Core Elements 2025 <https://www.cdc.gov/patient-safety/hcp/hospital-dx-excellence/index.html>
- Many SHEA White Papers 2024-2025

Predictive Modeling for Support of Empiric Choice at Time of Ordering

- INPIRE Trials. Gohil et al.
 - Pneumonia -- *JAMA*. 2024;331(23):2007-2017. doi:10.1001/jama.2024.6248
 - UTI -- *JAMA*. 2024;331(23):2018-2028. doi:10.1001/jama.2024.6259
 - IAI and SSTI -- *Open Forum Infect Dis*. 2025;12(Supplement_1):ofae631.059. doi:10.1093/ofid/ofae631.059

Data Modernization + AI Integration in Electronic Health Records

- Potential benefit in efficiency of AS work, but proof still in-process + barrier of investment resources + use while knowing limitations of AI...
- Data Sharing: <https://www.cdc.gov/data-modernization/php/about/index.html>



58

A case, continued.

86F with history of dementia, diabetes, and poor functional status presents from SNF with confusion, fever.

Day 3: Remains on vancomycin and zosyn. Progress note still says "sepsis." BCx negative. Awake/interactive.


Clinical pharmacist reviews the patient for vancomycin dosing, sees urine +E. coli susceptible to multiple oral and intravenous agents.

Contacts the provider.



59

Clinical Discussion



Tamma PD et al. *JAMA*. 2019;321(2):139-140.

600

60

Summary

Antibiotics are life-saving medicines that are often misused.

Antimicrobial decision-making is complex.

Optimized antimicrobial use through antimicrobial stewardship protects patients from unintended consequences.

Antimicrobial use affects individuals AND populations. Healthcare exposed populations are the most at risk.

Antimicrobial Stewardship Programs are required in US healthcare facilities and a key component of infection prevention.



61

THANK YOU!

Rebekah.Moehring@duke.edu



dcasip.medicine.duke.edu



62

CONTENT CUT FROM TALK

63

Antibiotics with Gram Positive (+) Activity

<i>S. aureus</i>	MRSA	VRE	<i>E. faecalis</i>
Nafcillin/Oxacillin			Ampicillin
Ampicillin/Sulbactam, Piperacillin/Tazobactam			Ampicillin/Sulbactam, Piperacillin/Tazobactam
Cephalosporins	Ceftaroline (only)		
Carbapenems			
(Fluoroquinolones)			
Vancomycin	Vancomycin		Vancomycin
Clindamycin	Clindamycin +/-		
Linezolid	Linezolid	Linezolid	Linezolid
Daptomycin	Daptomycin	Daptomycin	Daptomycin
Telavancin	Telavancin		
TMP-SMX	TMP-SMX		
Dalvabancin, Oritavancin	Dalvabancin, Oritavancin		

64

Antibiotics with Gram Negative (-) Activity

<i>E. coli</i>	<i>K. pneumoniae</i>	<i>Enterobacter</i>	<i>P. aeruginosa</i>
(Ampicillin)			
(Amp/sulb)	(Amp/sulb)		
Pip/Tazo	Pip/Tazo	Pip/Tazo	Pip/Tazo
Cephalosporins	Cephalosporins	3 rd , 4 th , 5 th gen.	Ceftaz/Cefepime
Carbapenems	Carbapenems	Carbapenems	Imip, Mero, Dori
Aztreonam	Aztreonam	Aztreonam	Aztreonam
Aminoglycosides	Aminoglycosides	Aminoglycosides	Amino-glycosides
Fluoroquinolone	Fluoroquinolone	Fluoroquinolone	Cipro and Levo
Trimeth/Sulf	Trimeth/Sulf	Trimeth/Sulf	



65

Antibiotic use assoc. with *CDI* risk – even in patients who don't get Abx!


46-month, single center retrospective study

255 of 34,298 patients developed CDI
(Incidence rate, 5.95 per 10,000 ptd)

Each 10% increase in ward-level antibiotic exposure was associated with a 2.1 per 10,000 increase in CDI

After adjustment for patient-level RFs, effect persisted: Relative risk, 1.34 (1.16-1.57) per 10% increase in days of therapy

Figure 2. Ward *Clostridium difficile* Infection (CDI) Incidence and Antibiotic Use Across Hospital Wards and Among Patients With and Without Direct Antibiotic Exposure



Each pair of numbered symbols represents the incidence of *C difficile* infection among the subset of patients who received antibiotics (diamonds) and those who did not (circles) within a given ward. For correspondence of ward identifiers, see Table 2. DOTs indicates days of therapy.




JAMA Intern Med. 2015;175(4):626-633

66

GRAM POSITIVE ORGANISMS

Gram positive cocci


- *Staphylococcus aureus*
- Coagulase negative staphylococcus
- *Streptococcus pneumoniae*
- *Streptococcus* sp.
- *Enterococcus* sp.



Gram positive – skin, lung, guts, devices

Gram positive rods

- *Bacillus* sp. (aerobes)
- *Clostridium* sp. (anaerobes)



MSSA:
Cefazolin
Nafcillin/Oxacillin

MRSA:
Vancomycin
Daptomycin
TMP/SMX
Linezolid

Strep:
Penicillin
Ampicillin/Amoxicillin

Ceftriaxone

Enterococci: Ampicillin,
Vancomycin

VRE: Daptomycin, Linezolid




67

GRAM NEGATIVE ORGANISMS

Gram negative cocci

- *Neisseria meningitidis*
- *Neisseria gonorrhoeae*




Gram negative rods (enteric)

- *E. coli*
- *Klebsiella* sp.
- *Enterobacter* sp.
- *Proteus* sp.
- *Serratia* sp.

Gram negative rods (non-enteric, non-lactose fermenters)

- *Pseudomonas aeruginosa*
- *Stenotrophomonas maltophilia*
- *Acinetobacter* sp.



CA-Enterics:
Ceftriaxone
Quinolones
TMP/SMX


Gram- negative: guts, urine, some lung

Anti-pseudomonal or HA-Resistant Enterics:
Pip/tazo
Cefepime
Ceftazidime
(Carbapenem)



68

Antibiotics with Anti-anaerobic Activity



Class	Agents (Route)	<i>B. fragilis</i> susceptibility ⁴⁻⁷
Beta-lactam beta-lactamase inhibitor combinations	amoxicillin/clav (PO)	90-97%
	ampicillin/sulb (IV)	97%
	piperacillin/tazo (IV)	> 99%
Cephalosporin	cefotetan (IV)	N/A
	cefoxitin (IV)	83-90%
Carbapenem	doripenem (IV)	> 99%
	ertapenem (IV/IM)	
	meropenem (IV)	
	imipenem (IV)	
Fluoroquinolone	moxifloxacin (IV/PO)	66-70%
Other	clindamycin (IV/PO)	66-70%
	metronidazole (IV/PO)	> 99%
	tigecycline (IV)	81-96%

B. fragilis is the most common group of gut anaerobes.

Then GPRs (*Clostridium* spp.)

Also consider: mouth, vaginal sources

Gas- and abscess-forming, bad odors


C. difficile is a special case (oral vancomycin).



69

NON-STAINING Bacterial PATHOGENS

- “Atypicals” -- Not stained by Gram’s method (Intracellular)
 - Legionella* sp. (Antigen test of the urine)
 - Chlamydia* (PCR)
 - Rickettsia* (Serology or PCR)
- Mycobacteria
 - M. tuberculosis*
 - Non-tuberculous mycobacteria



Ziehl-Neelsen Stain of TB

Macrolides:
Azithromycin
Clarithromycin

Tetracyclines:
Doxycycline
Minocycline

Community-acquired pneumonia	Pathogens	CXR pattern
Typical pneumonia	Bacterial: <i>S. Pneumoniae</i> <i>H. Influenzae</i>	Lobar, dense
Atypical pneumonia	Viral: influenza, RSV	Diffuse, patchy
	Bacterial: <i>Legionella</i> <i>Mycoplasma</i> <i>Chlamydia</i>	



70

Fungi

Candida/Yeast

- Fluconazole
- Echinocandins (micafungin, caspofungin)


Cryptococcus

- Amphotericin + flucytosine
- Fluconazole

Molds (Rhizopus/Mucor, Aspergillus)

- Voriconazole
- Posaconazole, Isavuconazole
- Itraconazole

<http://drfungus.org>



Guts, devices,
immunosuppressed
+ abx-exposed
hosts



71

DIAGNOSIS: Antigen Tests


- Very useful for following (and sometimes diagnosing) viral infections: HIV, HBV, COVID-19
- Occasionally useful for other pathogens (e.g., Cryptococcus)
- Quick/rapid tests are often based on antigen testing, less expensive
- Urine is sometimes used for antigen tests (e.g. Legionella urine antigen)
- Limitations: Less Sensitivity



72

DIAGNOSIS: Serology or Antibody Tests

- For bacterial infections, generally not useful in early diagnosis (usually requires acute and convalescent tests)
- For viral infections, IgM may allow early diagnosis (e.g., HepA)
- Works ok for difficult to access/culture pathogens
- Limitation: slower turnaround, cannot distinguish phase of illness
 - E.g. Syphilis IgG -- after initial infection, test remains positive for lifetime




73

DIAGNOSIS: PCR and Molecular tests

- Highly sensitive
- Allows diagnosis of non-culturable pathogens (e.g., norovirus)
- Limitations: Subject to false positives (e.g. *C. difficile* colonization vs. infection?)
- Rapid turnaround time
- Increasing use of Syndromic Panel tests (e.g. blood, respiratory, CSF, GI) that also include resistance genes
- E.g. Blood culture + GNR – get pathogen ID and molecular resistance gene targets within 1-2 hours



74



75

MIC = Minimal inhibitory concentration. Lowest concentration of antimicrobial that inhibits growth of bacteria. Commonly used in clinical lab.

Breakpoint = The MIC that is used to designate between susceptible and resistant.

Intrinsic vs. Acquired

Known quantity of bacteria placed into each tube

0.25 µg/mL 0.5 µg/mL 1.0 µg/mL 2.0 µg/mL **4.0 µg/mL** 8.0 µg/mL 16 µg/mL

Increasing antibiotic concentration →

Duke Center for Antimicrobial Stewardship and Infection Prevention

76

Resources for LTC Stewardship



JAMDA
journal homepage: www.jamda.com

Special Article

Template for an Antibiotic Stewardship Policy for Post-Acute and Long-Term Care Settings

Robin L.P. Jump MD, PhD^{a,b,*}, Swati Gaur MD, MBA, CMD^c, Morgan J. Katz MD^d, Christopher J. Crnich MD, PhD^{e,f}, Ghinwa Dumyati MD^g, Muhammad S. Ashraf MBBS^h, Elizabeth Frentzel MPHⁱ, Steven J. Schweon RN, MPH, MSN, CIC, HEM^j, Philip Sloane MD, MPH^k, David Nace MD, MPH, CMD^l on behalf of the Infection Advisory Committee for AMDA—The Society of Post-Acute and Long-Term Care Medicine

- CDC Core Elements
- AHRQ Guide
- CMS Standard Interpretive Guidance



Table 1

Resources for Antibiotic Stewardship in LTC

Category	Institution	Resource (type of link)
General Antibiotic Stewardship Principles	CDC*	The Core Elements of Antibiotic Stewardship for Nursing Homes (website) Core Elements of Antibiotic Stewardship for Nursing Homes (pdf) Checklist for the Core Elements of Antibiotic Stewardship for Nursing Homes (pdf) Nursing Home Antimicrobial Stewardship Guide (website) Do Bugs Need Drugs? (website) Rochester Nursing Home Collaborative (website)
	AHRQ	Minnesota Antimicrobial Stewardship Program Toolkit for Long-term Care Facilities (website) Antimicrobial Stewardship in Long-Term Care (pdf) Evaluation & Treatment –UTI in Elderly (website)
	AHS and BC CDC	Promoting Wise Antibiotic Use in Nursing Homes (pdf) Policy and Practice Actions to Improve Antibiotic Use (pdf)
	Rochester Patient Safety C. difficile Prevention Collaborative	Leading Antibiotic Stewardship in Nursing Homes (pdf) Creating a Culture to Improve Antibiotic Use in Nursing Homes (pdf) Starting an Antimicrobial Stewardship Program (website)
Policy and Implementation	Minnesota Department of Health	Monitor and Sustain Stewardship (website) Action Steps and Strategies for Implementing Antimicrobial Stewardship in Long-term Care Facilities (pdf)
	Massachusetts Coalition University of North Carolina CDC	Antimicrobial Stewardship Gap Analysis Tool (pdf) The Core Elements of Antibiotic Stewardship with CMS and QAPI Updates (pdf) Common Suspected Infections: Communication and Decision Making for Four Infections (website) Suspected UTI/SIAB Toolkit (website)
Antibiotic Use Protocols	New York Department of Health	Minimum Criteria for Common Infections Toolkit (website) Managing Common Infections in Older Adults (pdf) Guidelines for Treatment of Urinary Tract Infections (pdf)
	AHRQ	Minimum Criteria for Initiation of Antibiotics in Long-Term Care Residents (pdf) ABCs for Diagnosing Urinary Tract Infection in Long-Term Care (pdf) Measures of Antibiotic Prescribing, Use and Outcomes (pdf)
Measuring and Monitoring Antibiotic Use	Rochester Patient Safety C. difficile Prevention Collaborative	Working With Your Lab to Improve Antibiotic Prescribing (website) Using Nursing Home Antibiograms to Choose the Right Antibiotic (website) Antibiotic Tracking Worksheet (excel file) Antibiotic Tracking Sheet Instructions for Use (word document) Antibiotic Order Sheet Template (pdf) Antimicrobial Use Assessment for Long-term Care Facilities (pdf) About Antibiograms (pdf)
	Minnesota Department of Health	WHO/NET Collaborating Centre for Surveillance of Antimicrobial Resistance (Website) What to Ask Your Healthcare Provider about Antibiotics (pdf) What You Need to Know About Antibiotics in a Nursing Home (pdf) Toolkit to Educate and Engage Residents and Family Members (website)
Family and Resident Education	World Health Organization	Be Smart About Antibiotics (pdf) Talking With Residents (pdf) Talking With Residents' Family Members (pdf) Resident Information Sheet: Antibiotic-Resistant Bacteria (pdf) AMDA Choosing Wisely List (pdf) ACS Choosing Wisely List (website with pdfs)
	CDC	Track & Treatments for UTIs in older people – When you need them and when you don't (website with pdfs) FAQ for Families, Guardians and Health Care Aides-UTIs in LTCF (pdf) FAQ for Families, Guardians and Health Care Aides-NHAP in LTCF (pdf) Antibiotics for Urinary Tract Infections in Older Adults (pdf) Appropriate Bacteriuria Family Letter Template (pdf) Why Not Antibiotics? (pdf)
	AHRQ	Why Not Antibiotics? (pdf)
	ABIM	
	AHS and BC CDC	
	Rochester Patient Safety C. difficile Prevention Collaborative	
	University of North Carolina	

ABIM, American Board of Internal Medicine; ACS, American Geriatrics Society; AHRQ, Agency for Healthcare Research and Quality; AHS, Alberta Health Services; BC CDC, British Columbia Center for Disease Control; CDC, Centers for Disease Control and Prevention; UTI, urinary tract infection.

CDC: What to Do? 2019



Table. The Core Elements of Hospital Antibiotic Stewardship for Hospital Core Element Implementation.

Hospitals that have implemented the Hospital Core Elements of Antibiotic Stewardship Implementation to further enhance their stewardship program.

Hospital Core Elements	
Hospital Leadership Commitment	
	Dedicate necessary human, financial, and information technology resources.
Accountability	
	Appoint a leader or co-leaders, such as a physician and pharmacist, responsible for program management and outcomes.
Pharmacy/Stewardship Expertise	
	Appoint a pharmacist, ideally as the co-leader of the stewardship program, to help lead implementation efforts to improve antibiotic use.
Action	
	Implement interventions, such as prospective audit and feedback or preauthorization, to improve antibiotic use.
Tracking	
	Monitor antibiotic prescribing, impact of interventions, and other important outcomes, like C. difficile infections and resistance patterns.
Reporting	
	Regularly report information on antibiotic use and resistance to prescribers, pharmacists, nurses, and hospital leadership.
Education	
	Educate prescribers, pharmacists, nurses, and patients about adverse reactions from antibiotics, antibiotic resistance, and optimal prescribing.

* For critical access hospitals (CAHs), this criterion can be met if the hospital involved in stewardship (recognizing that some CAHs do not have pharmacy services).

1. Dedicate FTE
2. Accountable leaders
3. Pharmacist
4. Act: Do something
5. Track: AU Rates + resistance + HAIs
6. Share data back
7. Do some education